

CXCR5 Antibody
Rabbit Polyclonal Antibody
Catalog # ALS12496**Specification**

CXCR5 Antibody - Product Information

| | |
|-------------------|------------------------|
| Application | IHC |
| Primary Accession | P32302 |
| Reactivity | Human |
| Host | Rabbit |
| Clonality | Polyclonal |
| Calculated MW | 42kDa KDa |

CXCR5 Antibody - Additional Information**Gene ID** 643**Other Names**

C-X-C chemokine receptor type 5, CXC-R5, CXCR-5, Burkitt lymphoma receptor 1, Monocyte-derived receptor 15, MDR-15, CD185, CXCR5, BLR1, MDR15

Target/Specificity

The antibody recognizes 36 kD CXCR5 of human origin. Reactivity to other species has not been tested.

Reconstitution & Storage

Long term: -70°C; Short term: -20°C; Avoid freeze-thaw cycles.

Precautions

CXCR5 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

CXCR5 Antibody - Protein Information**Name** CXCR5**Synonyms** BLR1, MDR15**Function**

Cytokine receptor that binds to B-lymphocyte chemoattractant (BLC). Involved in B-cell migration into B-cell follicles of spleen and Peyer patches but not into those of mesenteric or peripheral lymph nodes. May have a regulatory function in Burkitt lymphoma (BL) lymphomagenesis and/or B-cell differentiation.

Cellular Location

Cell membrane; Multi-pass membrane protein.

Tissue Location


Expression in mature B-cells and Burkitt lymphoma cells

CXCR5 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CXCR5 Antibody - Images



Anti-CXCR5 antibody IHC of human spleen.

CXCR5 Antibody - Background

Cytokine receptor that binds to B-lymphocyte chemoattractant (BLC). Involved in B-cell migration into B-cell follicles of spleen and Peyer patches but not into those of mesenteric or peripheral lymph nodes. May have a regulatory function in Burkitt lymphoma (BL) lymphomagenesis and/or B-cell differentiation.

CXCR5 Antibody - References

- Dobner T., et al. Eur. J. Immunol. 22:2795-2799(1992).
Barella L., et al. Biochem. J. 309:773-779(1995).
Legler D.F., et al. J. Exp. Med. 187:655-660(1998).
Sjoebloom T., et al. Science 314:268-274(2006).