

F7 / Factor VII Antibody
Sheep Polyclonal Antibody
Catalog # ALS12590

Specification

F7 / Factor VII Antibody - Product Information

Application	IHC
Primary Accession	P08709
Reactivity	Human
Host	Sheep
Clonality	Polyclonal
Calculated MW	52kDa KDa

F7 / Factor VII Antibody - Additional Information

Gene ID 2155

Other Names

Coagulation factor VII, 3.4.21.21, Proconvertin, Serum prothrombin conversion accelerator, SPCA, Eptacog alfa, Factor VII light chain, Factor VII heavy chain, F7

Target/Specificity

It is specific for F. VII as determined by immunodiffusion. Identity was established with purified F. VII.

Reconstitution & Storage

Long term: -20°C; Short term: +4°C. Avoid repeat freeze-thaw cycles.

Precautions

F7 / Factor VII Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

F7 / Factor VII Antibody - Protein Information

Name F7

Function

Initiates the extrinsic pathway of blood coagulation. Serine protease that circulates in the blood in a zymogen form. Factor VII is converted to factor VIIa by factor Xa, factor XIIIa, factor IXa, or thrombin by minor proteolysis. In the presence of tissue factor and calcium ions, factor VIIa then converts factor X to factor Xa by limited proteolysis. Factor VIIa will also convert factor IX to factor IXa in the presence of tissue factor and calcium.

Cellular Location

Secreted.

Tissue Location


Plasma.

F7 / Factor VII Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

F7 / Factor VII Antibody - Images



Anti-F7 / Factor VII antibody IHC of human liver.

F7 / Factor VII Antibody - Background

Initiates the extrinsic pathway of blood coagulation. Serine protease that circulates in the blood in a zymogen form. Factor VII is converted to factor VIIa by factor Xa, factor XIIa, factor IXa, or thrombin by minor proteolysis. In the presence of tissue factor and calcium ions, factor VIIa then converts factor X to factor Xa by limited proteolysis. Factor VIIa will also convert factor IX to factor IXa in the presence of tissue factor and calcium.

F7 / Factor VII Antibody - References

- Hagen F.S., et al. Proc. Natl. Acad. Sci. U.S.A. 83:2412-2416(1986).
O'Hara P.J., et al. Proc. Natl. Acad. Sci. U.S.A. 84:5158-5162(1987).
Sabater-Lleal M., et al. Hum. Genet. 118:741-751(2006).
Soria J.M., et al. Submitted (DEC-2002) to the EMBL/GenBank/DDBJ databases.
Masroori N., et al. Submitted (MAR-2008) to the EMBL/GenBank/DDBJ databases.