

**PCK1 Antibody (Internal)**  
**Goat Polyclonal Antibody**  
**Catalog # ALS12627**

**Specification**

---

**PCK1 Antibody (Internal) - Product Information**

Application	IHC
Primary Accession	<a href="#">P35558</a>
Reactivity	Human, Rabbit, Monkey, Horse, Dog
Host	Goat
Clonality	Polyclonal
Calculated MW	69kDa KDa

**PCK1 Antibody (Internal) - Additional Information**

**Gene ID** 5105

**Other Names**

Phosphoenolpyruvate carboxykinase, cytosolic [GTP], PEPCK-C, 4.1.1.32, PCK1, PEPCK1

**Target/Specificity**

Human PCK1.

**Reconstitution & Storage**

Store at -20°C. Minimize freezing and thawing.

**Precautions**

PCK1 Antibody (Internal) is for research use only and not for use in diagnostic or therapeutic procedures.

**PCK1 Antibody (Internal) - Protein Information**

**Name** PCK1 {ECO:0000303|PubMed:8490617, ECO:0000312|HGNC:HGNC:8724}

**Function**

Cytosolic phosphoenolpyruvate carboxykinase that catalyzes the reversible decarboxylation and phosphorylation of oxaloacetate (OAA) and acts as the rate-limiting enzyme in gluconeogenesis (PubMed: [24863970](http://www.uniprot.org/citations/24863970), PubMed: [26971250](http://www.uniprot.org/citations/26971250), PubMed: [28216384](http://www.uniprot.org/citations/28216384), PubMed: [30193097](http://www.uniprot.org/citations/30193097)). Regulates cataplerosis and anaplerosis, the processes that control the levels of metabolic intermediates in the citric acid cycle (PubMed: [24863970](http://www.uniprot.org/citations/24863970), PubMed: [26971250](http://www.uniprot.org/citations/26971250), PubMed: [28216384](http://www.uniprot.org/citations/28216384), PubMed: [30193097](http://www.uniprot.org/citations/30193097)). At low glucose levels, it catalyzes the cataplerotic conversion of oxaloacetate to phosphoenolpyruvate

(PEP), the rate-limiting step in the metabolic pathway that produces glucose from lactate and other precursors derived from the citric acid cycle (PubMed:<a href="http://www.uniprot.org/citations/30193097" target="\_blank">30193097</a>). At high glucose levels, it catalyzes the anaplerotic conversion of phosphoenolpyruvate to oxaloacetate (PubMed:<a href="http://www.uniprot.org/citations/30193097" target="\_blank">30193097</a>). Acts as a regulator of formation and maintenance of memory CD8(+) T-cells: up- regulated in these cells, where it generates phosphoenolpyruvate, via gluconeogenesis (By similarity). The resultant phosphoenolpyruvate flows to glycogen and pentose phosphate pathway, which is essential for memory CD8(+) T-cells homeostasis (By similarity). In addition to the phosphoenolpyruvate carboxykinase activity, also acts as a protein kinase when phosphorylated at Ser-90: phosphorylation at Ser-90 by AKT1 reduces the binding affinity to oxaloacetate and promotes an atypical serine protein kinase activity using GTP as donor (PubMed:<a href="http://www.uniprot.org/citations/32322062" target="\_blank">32322062</a>). The protein kinase activity regulates lipogenesis: upon phosphorylation at Ser-90, translocates to the endoplasmic reticulum and catalyzes phosphorylation of INSIG proteins (INSIG1 and INSIG2), thereby disrupting the interaction between INSIG proteins and SCAP and promoting nuclear translocation of SREBP proteins (SREBF1/SREBP1 or SREBF2/SREBP2) and subsequent transcription of downstream lipogenesis- related genes (PubMed:<a href="http://www.uniprot.org/citations/32322062" target="\_blank">32322062</a>).

#### Cellular Location

Cytoplasm, cytosol. Endoplasmic reticulum Note=Phosphorylation at Ser-90 promotes translocation to the endoplasmic reticulum.

#### Tissue Location

Major sites of expression are liver, kidney and adipocytes.


### PCK1 Antibody (Internal) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### PCK1 Antibody (Internal) - Images





Anti-PCK1 antibody IHC of human liver.

### **PCK1 Antibody (Internal) - Background**

Catalyzes the conversion of oxaloacetate (OAA) to phosphoenolpyruvate (PEP), the rate-limiting step in the metabolic pathway that produces glucose from lactate and other precursors derived from the citric acid cycle.

### **PCK1 Antibody (Internal) - References**

- Stoffel M.,et al.Hum. Mol. Genet. 2:1-4(1993).
- Ting C.-N.,et al.Genomics 16:698-706(1993).
- Ota T.,et al.Nat. Genet. 36:40-45(2004).
- Deloukas P.,et al.Nature 414:865-871(2001).
- Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.