

**CD163 Antibody (clone GHI/61)**  
**Mouse Monoclonal Antibody**  
**Catalog # ALS12962**

**Specification**

---

**CD163 Antibody (clone GHI/61) - Product Information**

|                   |                        |
|-------------------|------------------------|
| Application       | IHC                    |
| Primary Accession | <a href="#">Q86VB7</a> |
| Reactivity        | Human                  |
| Host              | Mouse                  |
| Clonality         | Monoclonal             |
| Calculated MW     | 125kDa KDa             |

**CD163 Antibody (clone GHI/61) - Additional Information**

**Gene ID** 9332

**Other Names**

Scavenger receptor cysteine-rich type 1 protein M130, Hemoglobin scavenger receptor, CD163, Soluble CD163, sCD163, CD163, M130

**Target/Specificity**

CD163

**Reconstitution & Storage**

Store at 4°C, avoid repeated freeze thaw cycles.

**Precautions**

CD163 Antibody (clone GHI/61) is for research use only and not for use in diagnostic or therapeutic procedures.

**CD163 Antibody (clone GHI/61) - Protein Information**

**Name** CD163

**Synonyms** M130

**Function**

Acute phase-regulated receptor involved in clearance and endocytosis of hemoglobin/haptoglobin complexes by macrophages and may thereby protect tissues from free hemoglobin-mediated oxidative damage. May play a role in the uptake and recycling of iron, via endocytosis of hemoglobin/haptoglobin and subsequent breakdown of heme. Binds hemoglobin/haptoglobin complexes in a calcium-dependent and pH- dependent manner. Exhibits a higher affinity for complexes of hemoglobin and multimeric haptoglobin of HP\*1F phenotype than for complexes of hemoglobin and dimeric haptoglobin of HP\*1S phenotype. Induces a cascade of intracellular signals that involves tyrosine kinase-dependent calcium mobilization, inositol triphosphate production and secretion of IL6 and CSF1. Isoform 3 exhibits the higher capacity for ligand endocytosis and the more pronounced surface expression when expressed in cells.

### Cellular Location

[Soluble CD163]: Secreted

### Tissue Location


Expressed in monocytes and mature macrophages such as Kupffer cells in the liver, red pulp macrophages in the spleen, cortical macrophages in the thymus, resident bone marrow macrophages and meningeal macrophages of the central nervous system. Expressed also in blood. Isoform 1 is the lowest abundant in the blood. Isoform 2 is the lowest abundant in the liver and the spleen. Isoform 3 is the predominant isoform detected in the blood

### CD163 Antibody (clone GHI/61) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### CD163 Antibody (clone GHI/61) - Images



Anti-CD163 antibody IHC of human spleen.



Anti-CD163 antibody IHC of human liver.

### **CD163 Antibody (clone GHI/61) - Background**

Acute phase-regulated receptor involved in clearance and endocytosis of hemoglobin/haptoglobin complexes by macrophages and may thereby protect tissues from free hemoglobin-mediated oxidative damage. May play a role in the uptake and recycling of iron, via endocytosis of hemoglobin/haptoglobin and subsequent breakdown of heme. Binds hemoglobin/haptoglobin complexes in a calcium-dependent and pH-dependent manner. Exhibits a higher affinity for complexes of hemoglobin and multimeric haptoglobin of HP\*1F phenotype than for complexes of hemoglobin and dimeric haptoglobin of HP\*1S phenotype. Induces a cascade of intracellular signals that involves tyrosine kinase-dependent calcium mobilization, inositol triphosphate production and secretion of IL6 and CSF1. Isoform 3 exhibits the higher capacity for ligand endocytosis and the more pronounced surface expression when expressed in cells.

### **CD163 Antibody (clone GHI/61) - References**

- Law S.K.A., et al. Eur. J. Immunol. 23:2320-2325(1993).  
Ritter M., et al. Biochem. Biophys. Res. Commun. 260:466-474(1999).  
Welch S.-K.W., et al. Submitted (MAY-2005) to the EMBL/GenBank/DDBJ databases.  
Scherer S.E., et al. Nature 440:346-351(2006).  
Droste A., et al. Biochem. Biophys. Res. Commun. 256:110-113(1999).