

UCHL1 / PGP9.5 Antibody (aa171-220)
Rabbit Polyclonal Antibody
Catalog # ALS14295**Specification**

UCHL1 / PGP9.5 Antibody (aa171-220) - Product Information

Application	WB, IHC
Primary Accession	P09936
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	25kDa KDa

UCHL1 / PGP9.5 Antibody (aa171-220) - Additional Information**Gene ID** 7345**Other Names**

Ubiquitin carboxyl-terminal hydrolase isozyme L1, UCH-L1, 3.4.19.12, 6.-.-., Neuron cytoplasmic protein 9.5, PGP 9.5, PGP9.5, Ubiquitin thioesterase L1, UCHL1

Target/Specificity

PGP9.5 Antibody detects endogenous levels of total PGP9.5 protein.

Reconstitution & Storage

Store at -20°C.

Precautions

UCHL1 / PGP9.5 Antibody (aa171-220) is for research use only and not for use in diagnostic or therapeutic procedures.

UCHL1 / PGP9.5 Antibody (aa171-220) - Protein Information**Name** UCHL1**Function**

Deubiquitinase that plays a role in the regulation of several processes such as maintenance of synaptic function, cardiac function, inflammatory response or osteoclastogenesis (PubMed: [22212137](http://www.uniprot.org/citations/22212137), PubMed: [23359680](http://www.uniprot.org/citations/23359680)). Abrogates the ubiquitination of multiple proteins including WWTR1/TAZ, EGFR, HIF1A and beta-site amyloid precursor protein cleaving enzyme 1/BACE1 (PubMed: [22212137](http://www.uniprot.org/citations/22212137), PubMed: [25615526](http://www.uniprot.org/citations/25615526)). In addition, recognizes and hydrolyzes a peptide bond at the C-terminal glycine of ubiquitin to maintain a stable pool of monoubiquitin that is a key requirement for the ubiquitin-proteasome and the autophagy- lysosome pathways (PubMed: [12408865](http://www.uniprot.org/citations/12408865), PubMed: [8639624](http://www.uniprot.org/citations/8639624))

target="_blank">8639624, PubMed:9774100). Regulates amyloid precursor protein/APP processing by promoting BACE1 degradation resulting in decreased amyloid beta production (PubMed:22212137). Plays a role in the immune response by regulating the ability of MHC I molecules to reach cross-presentation compartments competent for generating Ag-MHC I complexes (By similarity). Mediates the 'Lys-48'-linked deubiquitination of the transcriptional coactivator WWTR1/TAZ leading to its stabilization and inhibition of osteoclastogenesis (By similarity). Deubiquitinates and stabilizes epidermal growth factor receptor EGFR to prevent its degradation and to activate its downstream mediators (By similarity). Modulates oxidative activity in skeletal muscle by regulating key mitochondrial oxidative proteins (By similarity). Enhances the activity of hypoxia-inducible factor 1-alpha/HIF1A by abrogating its VHL E3 ligase-mediated ubiquitination and consequently inhibiting its degradation (PubMed:25615526).

Cellular Location

Cytoplasm. Endoplasmic reticulum membrane; Lipid- anchor. Note=About 30% of total UCHL1 is associated with membranes in brain. Localizes near and/or within mitochondria to potentially interact with mitochondrial proteins {ECO:0000250|UniProtKB:Q9R0P9}

Tissue Location

Found in neuronal cell bodies and processes throughout the neocortex (at protein level). Expressed in neurons and cells of the diffuse neuroendocrine system and their tumors. Weakly expressed in ovary. Down-regulated in brains from Parkinson disease and Alzheimer disease patients.

Volume

50 µl


UCHL1 / PGP9.5 Antibody (aa171-220) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

UCHL1 / PGP9.5 Antibody (aa171-220) - Images






Western blot of extracts from 293 cells, using PGP9.5 Antibody.



Anti-UCHL1 / PGP9.5 antibody IHC of human brain, cortex.



Anti-UCHL1 / PGP9.5 antibody IHC of human kidney.

UCHL1 / PGP9.5 Antibody (aa171-220) - Background

Ubiquitin-protein hydrolase involved both in the processing of ubiquitin precursors and of ubiquitinated proteins. This enzyme is a thiol protease that recognizes and hydrolyzes a peptide bond at the C-terminal glycine of ubiquitin. Also binds to free monoubiquitin and may prevent its degradation in lysosomes. The homodimer may have ATP-independent ubiquitin ligase activity.

UCHL1 / PGP9.5 Antibody (aa171-220) - References

Hillier L.W., et al. Nature 434:724-731(2005).

Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Day I.N.M.,et al.Biochem. J. 268:521-524(1990).
Choi J.,et al.J. Biol. Chem. 279:13256-13264(2004).
Lubec G.,et al.Submitted (DEC-2008) to UniProtKB.