

CD74 Antibody (clone PIN.1)
Mouse Monoclonal Antibody
Catalog # ALS14971**Specification****CD74 Antibody (clone PIN.1) - Product Information**

Application	WB
Primary Accession	P04233
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Calculated MW	34kDa KDa

CD74 Antibody (clone PIN.1) - Additional Information**Gene ID** 972**Other Names**

HLA class II histocompatibility antigen gamma chain, HLA-DR antigens-associated invariant chain, Ia antigen-associated invariant chain, Ii, p33, CD74, CD74, DHLAG

Target/Specificity

Detects ~33-35 kD protein doublet, corresponding to the molecular mass of the p33 and p35 forms of human CD74 on SDS-PAGE immunoblots.

Reconstitution & Storage

Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles.

Precautions

CD74 Antibody (clone PIN.1) is for research use only and not for use in diagnostic or therapeutic procedures.

CD74 Antibody (clone PIN.1) - Protein Information**Name** CD74 ([HGNC:1697](#))**Synonyms** DHLAG**Function**

Plays a critical role in MHC class II antigen processing by stabilizing peptide-free class II alpha/beta heterodimers in a complex soon after their synthesis and directing transport of the complex from the endoplasmic reticulum to the endosomal/lysosomal system where the antigen processing and binding of antigenic peptides to MHC class II takes place. Serves as cell surface receptor for the cytokine MIF. [Isoform p41]: Stabilizes the conformation of mature CTSL by binding to its active site and serving as a chaperone to help maintain a pool of mature enzyme in endocytic compartments and extracellular space of antigen-presenting cells (APCs). Has antiviral activity by stymieing the endosomal entry of Ebola virus and coronaviruses, including SARS-CoV-2 (PubMed:32855215).

Disrupts cathepsin-mediated Ebola virus glycoprotein processing, which prevents viral fusion and entry. This antiviral activity is specific to p41 isoform (PubMed:32855215).

Cellular Location

Cell membrane; Single-pass type II membrane protein. Endoplasmic reticulum membrane. Golgi apparatus, trans-Golgi network. Endosome. Lysosome. Secreted. Note=Transits through a number of intracellular compartments in the endocytic pathway. It can either undergo proteolysis or reach the cell membrane

Tissue Location


Detected in urine (at protein level). [Isoform p33]: In B cells, represents 70% of total CD74 expression.

CD74 Antibody (clone PIN.1) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CD74 Antibody (clone PIN.1) - Images



CD-74 (PIN 1.1) N87 lysates mixed with macrophage inhibitory factor.

CD74 Antibody (clone PIN.1) - Background

Plays a critical role in MHC class II antigen processing by stabilizing peptide-free class II alpha/beta heterodimers in a complex soon after their synthesis and directing transport of the complex from the endoplasmic reticulum to the endosomal/lysosomal system where the antigen processing and binding of antigenic peptides to MHC class II takes place. Serves as cell surface receptor for the cytokine MIF.

CD74 Antibody (clone PIN.1) - References

- Claesson L., et al. Proc. Natl. Acad. Sci. U.S.A. 80:7395-7399(1983).
Strubin M., et al. EMBO J. 3:869-872(1984).
Kudo J., et al. Nucleic Acids Res. 13:8827-8841(1985).
O'Sullivan D.M., et al. Proc. Natl. Acad. Sci. U.S.A. 83:4484-4488(1986).
Kalinine N., et al. Submitted (OCT-2004) to the EMBL/GenBank/DDBJ databases.

