

**HNRNPD / AUF1 Antibody (aa49-98)**  
**Rabbit Polyclonal Antibody**  
**Catalog # ALS15261**

**Specification**

---

**HNRNPD / AUF1 Antibody (aa49-98) - Product Information**

Application	IF, WB, IHC
Primary Accession	<a href="#">O14103</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	38kDa KDa

**HNRNPD / AUF1 Antibody (aa49-98) - Additional Information**

**Gene ID** 3184

**Other Names**

Heterogeneous nuclear ribonucleoprotein D0, hnRNP D0, AU-rich element RNA-binding protein 1, HNRNPD, AUF1, HNRPD

**Target/Specificity**

hnRPD Antibody detects endogenous levels of total hnRPD protein.

**Reconstitution & Storage**

Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles.

**Precautions**

HNRNPD / AUF1 Antibody (aa49-98) is for research use only and not for use in diagnostic or therapeutic procedures.

**HNRNPD / AUF1 Antibody (aa49-98) - Protein Information**

**Name** HNRNPD

**Synonyms** AUF1, HNRPD

**Function**

Binds with high affinity to RNA molecules that contain AU- rich elements (AREs) found within the 3'-UTR of many proto-oncogenes and cytokine mRNAs. Also binds to double- and single-stranded DNA sequences in a specific manner and functions a transcription factor. Each of the RNA-binding domains specifically can bind solely to a single-stranded non-monotonous 5'-UUAG-3' sequence and also weaker to the single-stranded 5'-TTAGGG-3' telomeric DNA repeat. Binds RNA oligonucleotides with 5'-UUAGGG-3' repeats more tightly than the telomeric single-stranded DNA 5'-TTAGGG-3' repeats. Binding of RRM1 to DNA inhibits the formation of DNA quadruplex structure which may play a role in telomere elongation. May be involved in translationally coupled mRNA turnover. Implicated with other RNA-binding proteins in the cytoplasmic deadenylation/translational and decay interplay of the FOS mRNA mediated by the major

coding-region determinant of instability (mCRD) domain. May play a role in the regulation of the rhythmic expression of circadian clock core genes. Directly binds to the 3'UTR of CRY1 mRNA and induces CRY1 rhythmic translation. May also be involved in the regulation of PER2 translation.

#### Cellular Location


Nucleus. Cytoplasm. Note=Localized in cytoplasmic mRNP granules containing untranslated mRNAs. Component of ribonucleosomes. Cytoplasmic localization oscillates diurnally

#### HNRNPD / AUF1 Antibody (aa49-98) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


#### HNRNPD / AUF1 Antibody (aa49-98) - Images




Immunofluorescence of HeLa cells, using hnRNP D (Ab-83) Antibody.



Western blot of extracts from 293/Jurkat/3T3/COLO205 cells, using hnRNP D (Ab-83) Antibody.



Anti-HNRNPD / AUF1 antibody IHC of human small intestine.



Anti-HNRNPD / AUF1 antibody IHC of human tonsil.

### **HNRNPD / AUF1 Antibody (aa49-98) - Background**

Binds with high affinity to RNA molecules that contain AU-rich elements (AREs) found within the 3'-UTR of many proto-oncogenes and cytokine mRNAs. Also binds to double- and single-stranded DNA sequences in a specific manner and functions as a transcription factor. Each of the RNA-binding domains specifically can bind solely to a single-stranded non-monotonous 5'-UUAG-3' sequence and also weaker to the single-stranded 5'-TTAGGG-3' telomeric DNA repeat. Binds RNA oligonucleotides with 5'-UUAGGG-3' repeats more tightly than the telomeric single-stranded DNA 5'-TTAGGG-3' repeats. Binding of RRM1 to DNA inhibits the formation of DNA quadruplex structure which may play a role in telomere elongation. May be involved in translationally coupled mRNA turnover. Implicated with other RNA-binding proteins in the cytoplasmic deadenylation/translational and decay interplay of the FOS mRNA mediated by the major coding-region determinant of instability (mCRD) domain. May play a role in the regulation of the rhythmic expression of circadian clock core genes. Directly binds to the 3'UTR of CRY1 mRNA and induces CRY1 rhythmic translation. May also be involved in the regulation of PER2 translation.

### **HNRNPD / AUF1 Antibody (aa49-98) - References**

- Kajita Y., et al. *J. Biol. Chem.* 270:22167-22175(1995).
- Dempsey L.A., et al. *Genomics* 49:378-384(1998).
- Ota T., et al. *Nat. Genet.* 36:40-45(2004).
- Hillier L.W., et al. *Nature* 434:724-731(2005).
- Mural R.J., et al. Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.