

KRT18 / CK18 / Cytokeratin 18 Antibody (clone DC-10)
Mouse Monoclonal Antibody
Catalog # ALS15310

Specification

KRT18 / CK18 / Cytokeratin 18 Antibody (clone DC-10) - Product Information

Application	IHC
Primary Accession	P05783
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Calculated MW	48kDa KDa

KRT18 / CK18 / Cytokeratin 18 Antibody (clone DC-10) - Additional Information

Gene ID 3875

Other Names

Keratin, type I cytoskeletal 18, Cell proliferation-inducing gene 46 protein, Cytokeratin-18, CK-18, Keratin-18, K18, KRT18, CYK18

Target/Specificity

Reacts with the 45 kDa human keratin 18.

Reconstitution & Storage

Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles. Store undiluted.

Precautions

KRT18 / CK18 / Cytokeratin 18 Antibody (clone DC-10) is for research use only and not for use in diagnostic or therapeutic procedures.

KRT18 / CK18 / Cytokeratin 18 Antibody (clone DC-10) - Protein Information

Name KRT18

Synonyms CYK18

Function

Involved in the uptake of thrombin-antithrombin complexes by hepatic cells (By similarity). When phosphorylated, plays a role in filament reorganization. Involved in the delivery of mutated CFTR to the plasma membrane. Together with KRT8, is involved in interleukin-6 (IL-6)-mediated barrier protection.

Cellular Location

Nucleus matrix {ECO:0000250|UniProtKB:Q5BJY9}. Cytoplasm, perinuclear region. Nucleus, nucleolus. Cytoplasm {ECO:0000250|UniProtKB:Q5BJY9}

Tissue Location

Expressed in colon, placenta, liver and very weakly in exocervix. Increased expression observed in lymph nodes of breast carcinoma.


Volume

100 µl

KRT18 / CK18 / Cytokeratin 18 Antibody (clone DC-10) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

KRT18 / CK18 / Cytokeratin 18 Antibody (clone DC-10) - Images

Anti-Cytokeratin 18 / Keratin 18 antibody IHC of human prostate, epithelium.

KRT18 / CK18 / Cytokeratin 18 Antibody (clone DC-10) - Background

Involved in the uptake of thrombin-antithrombin complexes by hepatic cells (By similarity). When phosphorylated, plays a role in filament reorganization. Involved in the delivery of mutated CFTR to the plasma membrane. Together with KRT8, is involved in interleukin-6 (IL-6)-mediated barrier protection.

KRT18 / CK18 / Cytokeratin 18 Antibody (clone DC-10) - References

- Oshima R.G., et al. Differentiation 33:61-68(1986).
Kim J.W., et al. Submitted (SEP-2004) to the EMBL/GenBank/DDBJ databases.
Kalnina N., et al. Submitted (OCT-2004) to the EMBL/GenBank/DDBJ databases.
Suzuki Y., et al. Submitted (APR-2005) to the EMBL/GenBank/DDBJ databases.
Kulesh D.A., et al. Mol. Cell. Biol. 8:1540-1550(1988).