

STK39 / SPAK Antibody (Internal)
Goat Polyclonal Antibody
Catalog # ALS15367**Specification**

STK39 / SPAK Antibody (Internal) - Product Information

Application	WB, IHC
Primary Accession	O9UEW8
Reactivity	Human, Rat, Rabbit, Hamster, Monkey, Horse, Dog
Host	Goat
Clonality	Polyclonal
Calculated MW	59kDa KDa

STK39 / SPAK Antibody (Internal) - Additional Information**Gene ID** 27347**Other Names**

STE20/SPS1-related proline-alanine-rich protein kinase, Ste-20-related kinase, 2.7.11.1, DCHT, Serine/threonine-protein kinase 39, STK39, SPAK

Target/Specificity

Human STK39 / SPAK.

Reconstitution & Storage

Store at -20°C. Minimize freezing and thawing.

Precautions

STK39 / SPAK Antibody (Internal) is for research use only and not for use in diagnostic or therapeutic procedures.

STK39 / SPAK Antibody (Internal) - Protein Information**Name** STK39**Function**

Effector serine/threonine-protein kinase component of the WNK-SPAK/OSR1 kinase cascade, which is involved in various processes, such as ion transport, response to hypertonic stress and blood pressure (PubMed: [16669787](http://www.uniprot.org/citations/16669787), PubMed: [18270262](http://www.uniprot.org/citations/18270262), PubMed: [21321328](http://www.uniprot.org/citations/21321328), PubMed: [34289367](http://www.uniprot.org/citations/34289367)). Specifically recognizes and binds proteins with a RFXV motif (PubMed: [16669787](http://www.uniprot.org/citations/16669787), PubMed: [21321328](http://www.uniprot.org/citations/21321328)). Acts downstream of WNK kinases (WNK1, WNK2, WNK3 or WNK4): following activation by WNK kinases, catalyzes phosphorylation of ion cotransporters, such as SLC12A1/NKCC2,

SLC12A2/NKCC1, SLC12A3/NCC, SLC12A5/KCC2 or SLC12A6/KCC3, regulating their activity (PubMed:21321328). Mediates regulatory volume increase in response to hyperosmotic stress by catalyzing phosphorylation of ion cotransporters SLC12A1/NKCC2, SLC12A2/NKCC1 and SLC12A6/KCC3 downstream of WNK1 and WNK3 kinases (PubMed:12740379, PubMed:16669787, PubMed:21321328). Phosphorylation of Na-K-Cl cotransporters SLC12A2/NKCC1 and SLC12A2/NKCC1 promote their activation and ion influx; simultaneously, phosphorylation of K-Cl cotransporters SLC12A5/KCC2 and SLC12A6/KCC3 inhibit their activity, blocking ion efflux (PubMed:16669787, PubMed:19665974, PubMed:21321328). Acts as a regulator of NaCl reabsorption in the distal nephron by mediating phosphorylation and activation of the thiazide-sensitive Na-Cl cotransporter SLC12A3/NCC in distal convoluted tubule cells of kidney downstream of WNK4 (PubMed:18270262). Mediates the inhibition of SLC4A4, SLC26A6 as well as CFTR activities (By similarity). Phosphorylates RELT (By similarity).

Cellular Location

Cytoplasm. Nucleus. Note=Nucleus when caspase-cleaved.

Tissue Location


Predominantly expressed in brain and pancreas followed by heart, lung, kidney, skeletal muscle, liver, placenta and testis.

STK39 / SPAK Antibody (Internal) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

STK39 / SPAK Antibody (Internal) - Images



STK39 antibody (0.5 ug/ml) staining of Jurkat lysate (35 ug protein/ml in RIPA buffer).



Anti-STK39 / SPAK antibody IHC of human brain, cortex.

STK39 / SPAK Antibody (Internal) - Background

May act as a mediator of stress-activated signals.

STK39 / SPAK Antibody (Internal) - References

- Johnston A.M., et al. *Oncogene* 19:4290-4297(2000).
Corominas R., et al. *Nat. Commun.* 5:3650-3650(2014).
Melnick M.B., et al. Submitted (OCT-1997) to the EMBL/GenBank/DDBJ databases.
Hillier L.W., et al. *Nature* 434:724-731(2005).
Baytel D., et al. Submitted (AUG-1997) to the EMBL/GenBank/DDBJ databases.