

CD71 / Transferrin Receptor Antibody (aa15-64)
Rabbit Polyclonal Antibody
Catalog # ALS16056**Specification**

CD71 / Transferrin Receptor Antibody (aa15-64) - Product Information

Application	WB
Primary Accession	P02786
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	85kDa KDa

CD71 / Transferrin Receptor Antibody (aa15-64) - Additional Information**Gene ID** 7037**Other Names**

Transferrin receptor protein 1, TR, TfR, TfR1, Trfr, T9, p90, CD71, Transferrin receptor protein 1, serum form, sTfR, TFRC

Target/Specificity

CD71/TfR (Ab-24) Antibody detects endogenous levels of total CD71/TfR protein.

Reconstitution & Storage

Store at -20°C for up to one year.

Precautions

CD71 / Transferrin Receptor Antibody (aa15-64) is for research use only and not for use in diagnostic or therapeutic procedures.

CD71 / Transferrin Receptor Antibody (aa15-64) - Protein Information**Name** TFRC**Function**

Cellular uptake of iron occurs via receptor-mediated endocytosis of ligand-occupied transferrin receptor into specialized endosomes (PubMed:[26214738](http://www.uniprot.org/citations/26214738)). Endosomal acidification leads to iron release. The apotransferrin-receptor complex is then recycled to the cell surface with a return to neutral pH and the concomitant loss of affinity of apotransferrin for its receptor. Transferrin receptor is necessary for development of erythrocytes and the nervous system (By similarity). A second ligand, the hereditary hemochromatosis protein HFE, competes for binding with transferrin for an overlapping C-terminal binding site. Positively regulates T and B cell proliferation through iron uptake (PubMed:[26642240](http://www.uniprot.org/citations/26642240)). Acts as a lipid sensor that regulates mitochondrial fusion by regulating activation of the JNK pathway (PubMed:[26214738](http://www.uniprot.org/citations/26214738)).

When dietary levels of stearate (C18:0) are low, promotes activation of the JNK pathway, resulting in HUWE1- mediated ubiquitination and subsequent degradation of the mitofusin MFN2 and inhibition of mitochondrial fusion (PubMed:26214738). When dietary levels of stearate (C18:0) are high, TFRC stearylation inhibits activation of the JNK pathway and thus degradation of the mitofusin MFN2 (PubMed:26214738). Mediates uptake of NICOL1 into fibroblasts where it may regulate extracellular matrix production (By similarity).

Cellular Location

Cell membrane; Single-pass type II membrane protein Melanosome. Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV

Volume


50 µl

CD71 / Transferrin Receptor Antibody (aa15-64) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CD71 / Transferrin Receptor Antibody (aa15-64) - Images



Western blot of extracts from 293 cells, treated with PMA 125 ng/ml 30', using CD71/TfR (Ab-24)...

CD71 / Transferrin Receptor Antibody (aa15-64) - Background

Cellular uptake of iron occurs via receptor-mediated endocytosis of ligand-occupied transferrin receptor into specialized endosomes. Endosomal acidification leads to iron release. The apotransferrin-receptor complex is then recycled to the cell surface with a return to neutral pH and the concomitant loss of affinity of apotransferrin for its receptor. Transferrin receptor is necessary for development of erythrocytes and the nervous system (By similarity). A second ligand, the

hereditary hemochromatosis protein HFE, competes for binding with transferrin for an overlapping C-terminal binding site.

CD71 / Transferrin Receptor Antibody (aa15-64) - References

Schneider C., et al. Nature 311:675-678(1984).

McClelland A., et al. Cell 39:267-274(1984).

Evans P., et al. Gene 199:123-131(1997).

Wheeler D.L., et al. Thesis (1999), University of Iowa, United States.

Totoki Y., et al. Submitted (MAR-2005) to the EMBL/GenBank/DDBJ databases.