

**CDKN2A / p16INK4a Antibody (clone 5A8A4,3G8D12)**  
**Mouse Monoclonal Antibody**  
**Catalog # ALS16384**

**Specification**

---

**CDKN2A / p16INK4a Antibody (clone 5A8A4,3G8D12) - Product Information**

Application	IHC
Primary Accession	<a href="#">P42771</a>
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Calculated MW	17kDa KDa

**CDKN2A / p16INK4a Antibody (clone 5A8A4,3G8D12) - Additional Information**

**Gene ID** 1029

**Other Names**

Cyclin-dependent kinase inhibitor 2A, isoforms 1/2/3, Cyclin-dependent kinase 4 inhibitor A, CDK4I, Multiple tumor suppressor 1, MTS-1, p16-INK4a, p16-INK4, p16INK4A, CDKN2A, CDKN2, MTS1

**Target/Specificity**

Human CDKN2A / p16 INK4A

**Format**

Presentation PBS, 0.03% Sodium Azide

**Reconstitution & Storage**

Short term +4°C; Long term -20°C

**Precautions**

CDKN2A / p16INK4a Antibody (clone 5A8A4,3G8D12) is for research use only and not for use in diagnostic or therapeutic procedures.

**CDKN2A / p16INK4a Antibody (clone 5A8A4,3G8D12) - Protein Information**

**Name** CDKN2A ([HGNC:1787](#))

**Synonyms** CDKN2, MTS1

**Function**

Acts as a negative regulator of the proliferation of normal cells by interacting strongly with CDK4 and CDK6. This inhibits their ability to interact with cyclins D and to phosphorylate the retinoblastoma protein.

**Cellular Location**

Cytoplasm. Nucleus

**Tissue Location**

Widely expressed but not detected in brain or skeletal muscle. Isoform 3 is pancreas-specific


**Volume**

Array

**CDKN2A / p16INK4a Antibody (clone 5A8A4,3G8D12) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**CDKN2A / p16INK4a Antibody (clone 5A8A4,3G8D12) - Images**

Human Breast: Formalin-Fixed, Paraffin-Embedded (FFPE)

**CDKN2A / p16INK4a Antibody (clone 5A8A4,3G8D12) - Background**

Acts as a negative regulator of the proliferation of normal cells by interacting strongly with CDK4 and CDK6. This inhibits their ability to interact with cyclins D and to phosphorylate the retinoblastoma protein.

**CDKN2A / p16INK4a Antibody (clone 5A8A4,3G8D12) - References**

Serrano M., et al. Nature 366:704-707(1993).  
Robertson K.D., et al. Oncogene 18:3810-3820(1999).  
Kitagawa Y., et al. J. Biol. Chem. 277:46289-46297(2002).  
Lin Y.C., et al. Oncogene 26:7017-7027(2007).  
Humphray S.J., et al. Nature 429:369-374(2004).

**CDKN2A / p16INK4a Antibody (clone 5A8A4,3G8D12) - Citations**

- [Gut bacteria identified in colorectal cancer patients promote tumourigenesis via butyrate secretion](#)