

**UVRAG Antibody (C-Terminus)**  
**Rabbit Polyclonal Antibody**  
**Catalog # ALS16426****Specification**

---

**UVRAG Antibody (C-Terminus) - Product Information**

Application	IF
Primary Accession	<a href="#">O9P2Y5</a>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	78kDa KDa

**UVRAG Antibody (C-Terminus) - Additional Information****Gene ID** 7405**Other Names**

UV radiation resistance-associated gene protein, p63, UVRAG

**Target/Specificity**

UVRAG antibody is human and mouse reactive.

**Reconstitution & Storage**

Long term: -20°C; Short term: +4°C. Avoid repeat freeze-thaw cycles.

**Precautions**

UVRAG Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

**UVRAG Antibody (C-Terminus) - Protein Information****Name** UVRAG**Function**

Versatile protein that is involved in regulation of different cellular pathways implicated in membrane trafficking. Involved in regulation of the COPI-dependent retrograde transport from Golgi and the endoplasmic reticulum by associating with the NRZ complex; the function is dependent on its binding to phosphatidylinositol 3-phosphate (PtdIns(3)P) (PubMed: [16799551](http://www.uniprot.org/citations/16799551), PubMed: [18552835](http://www.uniprot.org/citations/18552835), PubMed: [20643123](http://www.uniprot.org/citations/20643123), PubMed: [24056303](http://www.uniprot.org/citations/24056303), PubMed: [28306502](http://www.uniprot.org/citations/28306502)). During autophagy acts as a regulatory subunit of the alternative PI3K complex II (PI3KC3-C2) that mediates formation of phosphatidylinositol 3-phosphate and is believed to be involved in maturation of autophagosomes and endocytosis. Activates lipid kinase activity of PIK3C3 (PubMed: [16799551](http://www.uniprot.org/citations/16799551)),

PubMed: <a href="http://www.uniprot.org/citations/20643123" target="\_blank">20643123</a>, PubMed: <a href="http://www.uniprot.org/citations/24056303" target="\_blank">24056303</a>, PubMed: <a href="http://www.uniprot.org/citations/28306502" target="\_blank">28306502</a>). Involved in the regulation of degradative endocytic trafficking and cytokinesis, and in regulation of ATG9A transport from the Golgi to the autophagosome; the functions seems to implicate its association with PI3KC3-C2 (PubMed: <a href="http://www.uniprot.org/citations/16799551" target="\_blank">16799551</a>, PubMed: <a href="http://www.uniprot.org/citations/20643123" target="\_blank">20643123</a>, PubMed: <a href="http://www.uniprot.org/citations/24056303" target="\_blank">24056303</a>). Involved in maturation of autophagosomes and degradative endocytic trafficking independently of BECN1 but depending on its association with a class C Vps complex (possibly the HOPS complex); the association is also proposed to promote autophagosome recruitment and activation of Rab7 and endosome-endosome fusion events (PubMed: <a href="http://www.uniprot.org/citations/18552835" target="\_blank">18552835</a>, PubMed: <a href="http://www.uniprot.org/citations/28306502" target="\_blank">28306502</a>). Enhances class C Vps complex (possibly HOPS complex) association with a SNARE complex and promotes fusogenic SNARE complex formation during late endocytic membrane fusion (PubMed: <a href="http://www.uniprot.org/citations/24550300" target="\_blank">24550300</a>). In case of negative- strand RNA virus infection is required for efficient virus entry, promotes endocytic transport of virions and is implicated in a VAMP8- specific fusogenic SNARE complex assembly (PubMed: <a href="http://www.uniprot.org/citations/24550300" target="\_blank">24550300</a>).

#### Cellular Location

Late endosome. Lysosome. Cytoplasmic vesicle, autophagosome. Early endosome. Endoplasmic reticulum. Midbody. Chromosome, centromere. Note=Colocalizes with RAB9-positive compartments involved in retrograde transport from late endosomes to trans-Golgi network. Colocalization with early endosomes is only partial (PubMed:24056303). Recruited to autophagosome following interaction with RUBCNL/PACER (PubMed:28306502)

#### Tissue Location

Highly expressed in brain, lung, kidney and liver.


#### UVRAG Antibody (C-Terminus) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### UVRAG Antibody (C-Terminus) - Images





Immunofluorescence of UVRAG in human lung tissue with UVRAG antibody at 20 ug/ml.

#### **UVRAG Antibody (C-Terminus) - References**

- Perelman B.,et al.Genomics 41:397-405(1997).
- Iida A.,et al.Hum. Genet. 106:277-287(2000).
- Ota T.,et al.Nat. Genet. 36:40-45(2004).
- Taylor T.D.,et al.Nature 440:497-500(2006).
- Itakura E.,et al.Mol. Biol. Cell 19:5360-5372(2008).