

**NOS2 / iNOS Antibody (clone K13-A)**  
**Rabbit Polyclonal Antibody**  
**Catalog # ALS16792****Specification**

---

**NOS2 / iNOS Antibody (clone K13-A) - Product Information**

Application	IHC
Primary Accession	<a href="#">P35228</a>
Other Accession	<a href="#">4843</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	131117

**NOS2 / iNOS Antibody (clone K13-A) - Additional Information****Gene ID** 4843**Other Names**

NOS2, Hepatocyte NOS, HEP-NOS, Inducible NOS, NOS, type II, NOS type II, Inducible NO synthase, NOS2A, INOS, NOS

**Reconstitution & Storage**20 mM Tris-HCl, pH 8.0, 20 mg/ml BSA stabilizer, 0.05% NaN<sub>3</sub> preservative. +4°C, avoid freezing**Precautions**

NOS2 / iNOS Antibody (clone K13-A) is for research use only and not for use in diagnostic or therapeutic procedures.

**NOS2 / iNOS Antibody (clone K13-A) - Protein Information****Name** NOS2 ([HGNC:7873](#))**Synonyms** NOS2A**Function**

Produces nitric oxide (NO) which is a messenger molecule with diverse functions throughout the body (PubMed:[7504305](http://www.uniprot.org/citations/7504305), PubMed:[7531687](http://www.uniprot.org/citations/7531687), PubMed:[7544004](http://www.uniprot.org/citations/7544004), PubMed:[7682706](http://www.uniprot.org/citations/7682706)). In macrophages, NO mediates tumoricidal and bactericidal actions. Also has nitrosylase activity and mediates cysteine S-nitrosylation of cytoplasmic target proteins such PTGS2/COX2 (By similarity). As component of the iNOS-S100A8/9 transnitrosylase complex involved in the selective inflammatory stimulus-dependent S-nitrosylation of GAPDH on 'Cys-247' implicated in regulation of the GAIT complex activity and probably multiple targets including ANXA5, EZR, MSN and VIM (PubMed:[25417112](http://www.uniprot.org/citations/25417112)). Involved in

inflammation, enhances the synthesis of pro-inflammatory mediators such as IL6 and IL8 (PubMed:<a href="http://www.uniprot.org/citations/19688109" target="\_blank">19688109</a>).

#### Cellular Location

Cytoplasm, cytosol. Note=Localizes as discrete foci scattered throughout the cytosol and in the presence of SPSB1 and SPSB4, exhibits a more diffuse cytosolic localization.

#### Tissue Location

Expressed in the liver, retina, bone cells and airway epithelial cells of the lung. Not expressed in the platelets Expressed in chondrocytes (PubMed:7504305)

#### Volume


50 µl

### NOS2 / iNOS Antibody (clone K13-A) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### NOS2 / iNOS Antibody (clone K13-A) - Images



Anti-NOS2 / iNOS antibody IHC staining of human liver.

### NOS2 / iNOS Antibody (clone K13-A) - Background

Produces nitric oxide (NO) which is a messenger molecule with diverse functions throughout the body. In macrophages, NO mediates tumoricidal and bactericidal actions. Also has nitrosylase activity and mediates cysteine S-nitrosylation of cytoplasmic target proteins such COX2.

### NOS2 / iNOS Antibody (clone K13-A) - References

Sherman P.A.,et al.Biochemistry 32:11600-11605(1993).  
Geller D.A.,et al.Proc. Natl. Acad. Sci. U.S.A. 90:3491-3495(1993).  
Charles I.G.,et al.Proc. Natl. Acad. Sci. U.S.A. 90:11419-11423(1993).  
Maier R.,et al.Biochim. Biophys. Acta 1208:145-150(1994).

Park C.S., et al. Biochem. Biophys. Res. Commun. 205:85-91(1994).