

VEGFA / VEGF Antibody (N-Terminus)
Rabbit Polyclonal Antibody
Catalog # ALS16808

Specification

VEGFA / VEGF Antibody (N-Terminus) - Product Information

Application	IHC
Primary Accession	P15692
Other Accession	7422
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	43597

VEGFA / VEGF Antibody (N-Terminus) - Additional Information

Gene ID 7422

Other Names

VEGFA, VPF, Vascular permeability factor, VEGF, VEGF-A, MVCD1

Target/Specificity

Human, mouse, rat. No cross reactivity with other proteins.

Reconstitution & Storage

5 mg BSA, 0.9 mg sodium chloride, 0.2 mg sodium phosphate, 0.05 mg Thimerosal, 0.05 mg sodium azide. Short term +4°C; Long term -20°C

Precautions

VEGFA / VEGF Antibody (N-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

VEGFA / VEGF Antibody (N-Terminus) - Protein Information

Name VEGFA

Synonyms VEGF

Function

[N-VEGF]: Participates in the induction of key genes involved in the response to hypoxia and in the induction of angiogenesis such as HIF1A (PubMed:35455969). Involved in protecting cells from hypoxia- mediated cell death (By similarity).

Cellular Location

[N-VEGF]: Cytoplasm. Nucleus. Note=Cytoplasmic in normoxic conditions and localizes to the nucleus under hypoxic conditions [Isoform L-VEGF189]: Endoplasmic reticulum. Golgi apparatus.

Secreted, extracellular space, extracellular matrix [Isoform VEGF165]: Secreted

Tissue Location

Higher expression in pituitary tumors than the pituitary gland. [Isoform VEGF165]: Widely expressed. [Isoform VEGF206]: Not widely expressed.

Volume


50 μ l

VEGFA / VEGF Antibody (N-Terminus) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

VEGFA / VEGF Antibody (N-Terminus) - Images



Anti-VEGFA / VEGF antibody IHC staining of human small intestine.

VEGFA / VEGF Antibody (N-Terminus) - Background

Growth factor active in angiogenesis, vasculogenesis and endothelial cell growth. Induces endothelial cell proliferation, promotes cell migration, inhibits apoptosis and induces permeabilization of blood vessels. Binds to the FLT1/VEGFR1 and KDR/VEGFR2 receptors, heparan sulfate and heparin. NRP1/Neuropilin-1 binds isoforms VEGF-165 and VEGF-145. Isoform VEGF165B binds to KDR but does not activate downstream signaling pathways, does not activate angiogenesis and inhibits tumor growth.

VEGFA / VEGF Antibody (N-Terminus) - References

- Leung D.W., et al. *Science* 246:1306-1309(1989).
Keck P.J., et al. *Science* 246:1309-1312(1989).
Tischer E., et al. *J. Biol. Chem.* 266:11947-11954(1991).
Houck K.A., et al. *Mol. Endocrinol.* 5:1806-1814(1991).
Weindel K., et al. *Biochem. Biophys. Res. Commun.* 183:1167-1174(1992).