

**Anti-IGF1 Antibody**  
**Rabbit Anti Human Polyclonal Antibody**  
**Catalog # ALS17732****Specification**

---

**Anti-IGF1 Antibody - Product Information**

Application	<b>WB, IHC-P, E, Neut</b>
Primary Accession	<a href="#">P05019</a>
Predicted	<b>Human</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Isotype	<b>IgG</b>
Calculated MW	<b>21841</b>

**Anti-IGF1 Antibody - Additional Information****Gene ID** 3479**Alias Symbol** **IGF1****Other Names**

IGF1, IGF-IA, IGF-IB, IGF1, IBP1, IGF-I, Insulin-like growth factor 1, Insulin-like growth factor I, Insulin-like growth factor IA, Insulin-like growth factor IB, Mechano growth factor, Somatomedin-C, IGF1A, MGF

**Target/Specificity**

Human IGF-I

**Reconstitution & Storage**

Immunoaffinity purified

**Precautions**

Anti-IGF1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**Anti-IGF1 Antibody - Protein Information****Name** IGF1 ([HGNC:5464](#))**Function**

The insulin-like growth factors, isolated from plasma, are structurally and functionally related to insulin but have a much higher growth-promoting activity. May be a physiological regulator of [1-14C]- 2-deoxy-D-glucose (2DG) transport and glycogen synthesis in osteoblasts. Stimulates glucose transport in bone-derived osteoblastic (PyMS) cells and is effective at much lower concentrations than insulin, not only regarding glycogen and DNA synthesis but also with regard to enhancing glucose uptake. May play a role in synapse maturation (PubMed:<a href="http://www.uniprot.org/citations/21076856" target="\_blank">21076856</a>, PubMed:<a href="http://www.uniprot.org/citations/24132240" target="\_blank">24132240</a>). Ca(2+)-dependent exocytosis of IGF1 is required for sensory perception of smell in the olfactory bulb (By similarity). Acts as a ligand for IGF1R. Binds to the alpha subunit of IGF1R, leading to the

activation of the intrinsic tyrosine kinase activity which autophosphorylates tyrosine residues in the beta subunit thus initiating a cascade of down-stream signaling events leading to activation of the PI3K-AKT/PKB and the Ras-MAPK pathways. Binds to integrins ITGAV:ITGB3 and ITGA6:ITGB4. Its binding to integrins and subsequent ternary complex formation with integrins and IGFR1 are essential for IGF1 signaling. Induces the phosphorylation and activation of IGFR1, MAPK3/ERK1, MAPK1/ERK2 and AKT1 (PubMed:<a href="http://www.uniprot.org/citations/19578119" target="\_blank">19578119</a>, PubMed:<a href="http://www.uniprot.org/citations/22351760" target="\_blank">22351760</a>, PubMed:<a href="http://www.uniprot.org/citations/23243309" target="\_blank">23243309</a>, PubMed:<a href="http://www.uniprot.org/citations/23696648" target="\_blank">23696648</a>). As part of the MAPK/ERK signaling pathway, acts as a negative regulator of apoptosis in cardiomyocytes via promotion of STUB1/CHIP-mediated ubiquitination and degradation of ICER-type isoforms of CREM (By similarity).

#### Cellular Location

Secreted {ECO:0000250|UniProtKB:P05017}.

#### Anti-IGF1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Anti-IGF1 Antibody - Images