

**Anti-CD63 Antibody (clone NKI/C-3)**  
**Mouse Anti Human Monoclonal Antibody**  
**Catalog # ALS17758**

**Specification**

---

**Anti-CD63 Antibody (clone NKI/C-3) - Product Information**

Application	IHC-P
Primary Accession	<a href="#">P08962</a>
Predicted	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1,k
Calculated MW	25637

**Anti-CD63 Antibody (clone NKI/C-3) - Additional Information**

**Gene ID** 967

**Alias Symbol** CD63

**Other Names**

CD63, CD63 molecule, LAMP-3, ME491, Granulophysin, MLA1, Tetraspanin-30, Tspan-30, CD63 antigen, OMA81H, TSPAN30

**Reconstitution & Storage**

Protein G purified

**Precautions**

Anti-CD63 Antibody (clone NKI/C-3) is for research use only and not for use in diagnostic or therapeutic procedures.

**Anti-CD63 Antibody (clone NKI/C-3) - Protein Information**

**Name** CD63

**Synonyms** MLA1, TSPAN30

**Function**

Functions as a cell surface receptor for TIMP1 and plays a role in the activation of cellular signaling cascades. Plays a role in the activation of ITGB1 and integrin signaling, leading to the activation of AKT, FAK/PTK2 and MAP kinases. Promotes cell survival, reorganization of the actin cytoskeleton, cell adhesion, spreading and migration, via its role in the activation of AKT and FAK/PTK2. Plays a role in VEGFA signaling via its role in regulating the internalization of KDR/VEGFR2. Plays a role in intracellular vesicular transport processes, and is required for normal trafficking of the PMEL luminal domain that is essential for the development and maturation of melanocytes. Plays a role in the adhesion of leukocytes onto endothelial cells via its role in the regulation of SELP trafficking. May play a role in mast cell degranulation in response to Ms4a2/FcεRI stimulation, but not in mast cell degranulation in response to other stimuli.

**Cellular Location**

Cell membrane; Multi-pass membrane protein. Lysosome membrane; Multi-pass membrane protein. Late endosome membrane; Multi-pass membrane protein. Endosome, multivesicular body. Melanosome. Secreted, extracellular exosome. Cell surface. Note=Also found in Weibel-Palade bodies of endothelial cells (PubMed:10793155). Located in platelet dense granules (PubMed:7682577). Detected in a subset of pre-melanosomes Detected on intraluminal vesicles (ILVs) within multivesicular bodies (PubMed:21962903).

**Tissue Location**

Detected in platelets (at protein level). Dysplastic nevi, radial growth phase primary melanomas, hematopoietic cells, tissue macrophages.

**Anti-CD63 Antibody (clone NKI/C-3) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Anti-CD63 Antibody (clone NKI/C-3) - Images**