

Anti-PCNA Antibody (clone PC10)
Mouse Anti Human Monoclonal Antibody
Catalog # ALS17848**Specification**

Anti-PCNA Antibody (clone PC10) - Product Information

Application	WB, IHC-P, IHC-F, IP, FC
Primary Accession	P12004
Predicted	Human, Mouse, Rat, Rabbit, Hamster, Chicken, Rhesus, Sheep, Xenopus, Horse, Dog, Cat
Host	Mouse
Clonality	Monoclonal
Isotype	IgG2a
Calculated MW	28769

Anti-PCNA Antibody (clone PC10) - Additional Information**Gene ID 5111**

Alias Symbol	PCNA
Other Names	
PCNA, Cyclin	

Target/Specificity

Recognizes the proliferating cell nuclear antigen, also known as PCNA or cyclin. PCNA is a 261 amino acid ~28 kD nuclear protein vital for cellular DNA synthesis at the replication fork through its interaction with. PCNA is the auxilliary protein for ...

Reconstitution & Storage

Purified

Precautions

Anti-PCNA Antibody (clone PC10) is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-PCNA Antibody (clone PC10) - Protein Information**Name PCNA****Function**

Auxiliary protein of DNA polymerase delta and epsilon, is involved in the control of eukaryotic DNA replication by increasing the polymerase's processibility during elongation of the leading strand (PubMed:35585232). Induces a robust stimulatory effect on the 3'-5' exonuclease and 3'-phosphodiesterase, but not apurinic-apyrimidinic (AP) endonuclease, APEX2 activities. Has to be loaded onto DNA in order to be able to stimulate APEX2. Plays a key role in DNA damage response (DDR) by being conveniently positioned at the replication fork to coordinate DNA replication with DNA repair and DNA damage

tolerance pathways (PubMed:24939902). Acts as a loading platform to recruit DDR proteins that allow completion of DNA replication after DNA damage and promote postreplication repair: Monoubiquitinated PCNA leads to recruitment of translesion (TLS) polymerases, while 'Lys-63'-linked polyubiquitination of PCNA is involved in error-free pathway and employs recombination mechanisms to synthesize across the lesion (PubMed:24695737).

Cellular Location

Nucleus. Note=Colocalizes with CREBBP, EP300 and POLD1 to sites of DNA damage (PubMed:24939902). Forms nuclear foci representing sites of ongoing DNA replication and vary in morphology and number during S phase (PubMed:15543136). Co-localizes with SMARCA5/SNF2H and BAZ1B/WSTF at replication foci during S phase (PubMed:15543136). Together with APEX2, is redistributed in discrete nuclear foci in presence of oxidative DNA damaging agents

Anti-PCNA Antibody (clone PC10) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-PCNA Antibody (clone PC10) - Images