

**Anti-APOC2 / Apolipoprotein C II Antibody**  
**Rabbit Anti Human Polyclonal Antibody**  
**Catalog # ALS18135****Specification**

---

**Anti-APOC2 / Apolipoprotein C II Antibody - Product Information**

Application	<b>WB, IHC-P</b>
Primary Accession	<a href="#">P02655</a>
Predicted	<b>Human, Mouse, Rat</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Calculated MW	<b>11284</b>

**Anti-APOC2 / Apolipoprotein C II Antibody - Additional Information****Gene ID** 344**Alias Symbol** **APOC2**  
**Other Names**  
APOC2, Apolipoprotein C-II, APO-CII, Apolipoprotein C2, APOC-II**Target/Specificity**  
Human APOC2 / Apolipoprotein C II**Reconstitution & Storage**  
Immunoaffinity purified**Precautions**  
Anti-APOC2 / Apolipoprotein C II Antibody is for research use only and not for use in diagnostic or therapeutic procedures.**Anti-APOC2 / Apolipoprotein C II Antibody - Protein Information****Name** APOC2**Synonyms** APC2**Function**

Component of chylomicrons, very low-density lipoproteins (VLDL), low-density lipoproteins (LDL), and high-density lipoproteins (HDL) in plasma. Plays an important role in lipoprotein metabolism as an activator of lipoprotein lipase. Both proapolipoprotein C-II and apolipoprotein C-II can activate lipoprotein lipase. In normolipidemic individuals, it is mainly distributed in the HDL, whereas in hypertriglyceridemic individuals, predominantly found in the VLDL and LDL.

**Cellular Location**  
Secreted.**Tissue Location**

Liver and intestine..

### **Anti-APOC2 / Apolipoprotein C II Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **Anti-APOC2 / Apolipoprotein C II Antibody - Images**