

Anti-CYP27 / CYP27A1 Antibody (aa101-150)
Rabbit Anti Human Polyclonal Antibody
Catalog # ALS18317**Specification****Anti-CYP27 / CYP27A1 Antibody (aa101-150) - Product Information**

Application	WB, IHC-P, E
Primary Accession	Q02318
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	60235

Anti-CYP27 / CYP27A1 Antibody (aa101-150) - Additional Information**Gene ID 1593**Alias Symbol **CYP27A1****Other Names**CYP27A1, Cytochrome P450 27, CTX, Cytochrome P-450C27/25, CP27, Vitamin D(3)
25-hydroxylase, CYP27, Sterol 27-hydroxylase**Target/Specificity**

Cytochrome P450 27A1 Antibody detects endogenous levels of total Cytochrome P450 27A1 protein.

Reconstitution & Storage

Immunoaffinity purified

Precautions

Anti-CYP27 / CYP27A1 Antibody (aa101-150) is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-CYP27 / CYP27A1 Antibody (aa101-150) - Protein Information**Name** CYP27A1 {ECO:0000303|PubMed:21411718, ECO:0000312|HGNC:HGNC:2605}**Function**Cytochrome P450 monooxygenase that catalyzes regio- and stereospecific hydroxylation of cholesterol and its derivatives. Hydroxylates (with R stereochemistry) the terminal methyl group of cholesterol side-chain in a three step reaction to yield at first a C26 alcohol, then a C26 aldehyde and finally a C26 acid (PubMed: [12077124](http://www.uniprot.org/citations/12077124) target="_blank">12077124, PubMed: [21411718](http://www.uniprot.org/citations/21411718) target="_blank">21411718, PubMed: [28190002](http://www.uniprot.org/citations/28190002) target="_blank">28190002, PubMed: [9660774](http://www.uniprot.org/citations/9660774) target="_blank">9660774). Regulates cholesterol homeostasis by catalyzing the conversion of excess cholesterol to bile acids via both the 'neutral' (classic) and the 'acid' (alternative)

pathways (PubMed:11412116, PubMed:1708392, PubMed:2019602, PubMed:7915755, PubMed:9186905, PubMed:9660774, PubMed:9790667). May also regulate cholesterol homeostasis via generation of active oxysterols, which act as ligands for NR1H2 and NR1H3 nuclear receptors, modulating the transcription of genes involved in lipid metabolism (PubMed:12077124, PubMed:9660774). Plays a role in cholestanol metabolism in the cerebellum. Similarly to cholesterol, hydroxylates cholestanol and may facilitate sterol diffusion through the blood-brain barrier to the systemic circulation for further degradation (PubMed:28190002). Also hydroxylates retinal 7- ketocholesterol, a noxious oxysterol with pro-inflammatory and pro- apoptotic effects, and may play a role in its elimination from the retinal pigment epithelium (PubMed:21411718). May play a redundant role in vitamin D biosynthesis. Catalyzes 25-hydroxylation of vitamin D3 that is required for its conversion to a functionally active form (PubMed:15465040).

Cellular Location

Mitochondrion inner membrane {ECO:0000250|UniProtKB:P17178}; Peripheral membrane protein {ECO:0000250|UniProtKB:P17178}. Note=Post-translationally targeted to mitochondria. All three of the receptor proteins in the TOM complex, TOMM70, TOMM20 and TOMM22 are required for the translocation across the mitochondrial outer membrane. After translocation into the matrix, associates with the inner membrane as a membrane extrinsic protein {ECO:0000250|UniProtKB:P17178}

Tissue Location

Expressed in the neural retina and underlying retinal pigment epithelium (at protein level) (PubMed:21411718) Expressed in the gray and white matter of cerebellum (at protein level) (PubMed:28190002).

Anti-CYP27 / CYP27A1 Antibody (aa101-150) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-CYP27 / CYP27A1 Antibody (aa101-150) - Images