

**Anti-HMG1 / HMGB1 Antibody (aa1-200, clone ABM24D3)**  
**Mouse Anti Human Monoclonal Antibody**  
**Catalog # ALS18385**

### Specification

---

#### Anti-HMG1 / HMGB1 Antibody (aa1-200, clone ABM24D3) - Product Information

Application	WB, IHC, IHC-P, FC
Primary Accession	<a href="#">P09429</a>
Predicted	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1,k
Calculated MW	24894

#### Anti-HMG1 / HMGB1 Antibody (aa1-200, clone ABM24D3) - Additional Information

Gene ID 3146

Alias Symbol **HMGB1**

#### Other Names

HMGB1, Amphoterin, High mobility group protein B1, High mobility group box 1, High mobility group protein 1, HMG1, HMG3, SBP-1, High-mobility group box 1, HMG-1

#### Reconstitution & Storage

Purified

#### Precautions

Anti-HMG1 / HMGB1 Antibody (aa1-200, clone ABM24D3) is for research use only and not for use in diagnostic or therapeutic procedures.

#### Anti-HMG1 / HMGB1 Antibody (aa1-200, clone ABM24D3) - Protein Information

Name HMGB1 ([HGNC:4983](#))

Synonyms HMG1

#### Function

Multifunctional redox sensitive protein with various roles in different cellular compartments. In the nucleus is one of the major chromatin-associated non-histone proteins and acts as a DNA chaperone involved in replication, transcription, chromatin remodeling, V(D)J recombination, DNA repair and genome stability (PubMed:<a href="http://www.uniprot.org/citations/33147444" target="\_blank">33147444</a>). Proposed to be an universal biosensor for nucleic acids. Promotes host inflammatory response to sterile and infectious signals and is involved in the coordination and integration of innate and adaptive immune responses. In the cytoplasm functions as a sensor and/or chaperone for immunogenic nucleic acids implicating the activation of TLR9-mediated immune responses, and mediates autophagy. Acts as a danger-associated molecular pattern (DAMP) molecule that amplifies immune responses during tissue injury (PubMed:<a href="http://www.uniprot.org/citations/27362237" target="\_blank">27362237</a>).

Released to the extracellular environment can bind DNA, nucleosomes, IL-1 beta, CXCL12, AGER isoform 2/sRAGE, lipopolysaccharide (LPS) and lipoteichoic acid (LTA), and activates cells through engagement of multiple surface receptors (PubMed:<a href="http://www.uniprot.org/citations/34743181" target="\_blank">34743181</a>). In the extracellular compartment fully reduced HMGB1 (released by necrosis) acts as a chemokine, disulfide HMGB1 (actively secreted) as a cytokine, and sulfonyl HMGB1 (released from apoptotic cells) promotes immunological tolerance (PubMed:<a href="http://www.uniprot.org/citations/23446148" target="\_blank">23446148</a>, PubMed:<a href="http://www.uniprot.org/citations/23519706" target="\_blank">23519706</a>, PubMed:<a href="http://www.uniprot.org/citations/23994764" target="\_blank">23994764</a>, PubMed:<a href="http://www.uniprot.org/citations/25048472" target="\_blank">25048472</a>). Has proangiogenic activity (By similarity). May be involved in platelet activation (By similarity). Binds to phosphatidylserine and phosphatidylethanolamide (By similarity). Bound to RAGE mediates signaling for neuronal outgrowth (By similarity). May play a role in accumulation of expanded polyglutamine (polyQ) proteins such as huntingtin (HTT) or TBP (PubMed:<a href="http://www.uniprot.org/citations/23303669" target="\_blank">23303669</a>, PubMed:<a href="http://www.uniprot.org/citations/25549101" target="\_blank">25549101</a>).

### Cellular Location

Nucleus. Chromosome {ECO:0000250|UniProtKB:P10103, ECO:0000250|UniProtKB:P63159, ECO:0000305}. Cytoplasm. Secreted {ECO:0000250|UniProtKB:P63158, ECO:0000269|PubMed:12231511, ECO:0000269|PubMed:14532127, ECO:0000269|PubMed:15944249, ECO:0000269|PubMed:19811284, ECO:0000269|PubMed:22869893, ECO:0000269|PubMed:33147444}. Cell membrane {ECO:0000250|UniProtKB:P63158, ECO:0000250|UniProtKB:P63159, ECO:0000269|PubMed:11154118}; Peripheral membrane protein {ECO:0000250|UniProtKB:P63158, ECO:0000250|UniProtKB:P63159, ECO:0000269|PubMed:11154118}; Extracellular side {ECO:0000250|UniProtKB:P63158, ECO:0000250|UniProtKB:P63159, ECO:0000269|PubMed:11154118}. Endosome {ECO:0000250|UniProtKB:P63158} Endoplasmic reticulum-Golgi intermediate compartment {ECO:0000250|UniProtKB:P63158}. Note=In basal state predominantly nuclear. Shuttles between the cytoplasm and the nucleus (PubMed:12231511, PubMed:17114460). Translocates from the nucleus to the cytoplasm upon autophagy stimulation (PubMed:20819940). Release from macrophages in the extracellular milieu requires the activation of NLR4 or NLRP3 inflammasomes (By similarity). Passively released to the extracellular milieu from necrotic cells by diffusion, involving the fully reduced HGMB1 which subsequently gets oxidized (PubMed:19811284) Also released from apoptotic cells (PubMed:16855214, PubMed:18631454) Active secretion from a variety of immune and non-immune cells such as macrophages, monocytes, neutrophils, dendritic cells and natural killer cells in response to various stimuli such as LPS and cytokines involves a nonconventional secretory process via secretory lysosomes (PubMed:12231511, PubMed:14532127, PubMed:15944249). Secreted by plasma cells in response to LPS (By similarity). Found on the surface of activated platelets (PubMed:11154118). An increased chromatin association is observed when associated with the adenovirus protein pVII (PubMed:27362237). {ECO:0000250|UniProtKB:P63158, ECO:0000269|PubMed:11154118, ECO:0000269|PubMed:12231511, ECO:0000269|PubMed:14532127, ECO:0000269|PubMed:15944249, ECO:0000269|PubMed:16855214, ECO:0000269|PubMed:17114460, ECO:0000269|PubMed:18631454, ECO:0000269|PubMed:19811284, ECO:0000269|PubMed:20819940, ECO:0000269|PubMed:27362237, ECO:0000305|PubMed:20123072}

### Tissue Location

Ubiquitous. Expressed in platelets (PubMed:11154118).

### Anti-HMG1 / HMGB1 Antibody (aa1-200, clone ABM24D3) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Anti-HMG1 / HMGB1 Antibody (aa1-200, clone ABM24D3) - Images**