

Phospho-mBad(Ser155) Antibody
Purified Phospho-specific Monoclonal Antibody (Mab)
Catalog # AM1029a

Specification

Phospho-mBad(Ser155) Antibody - Product Information

Application	WB,E
Primary Accession	Q61337
Other Accession	NP_031548
Reactivity	Human, Mouse
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse IgG1

Phospho-mBad(Ser155) Antibody - Additional Information

Gene ID 12015

Other Names

Bcl2-associated agonist of cell death, BAD, Bcl-2-binding component 6, Bcl-xL/Bcl-2-associated death promoter, Bcl2 antagonist of cell death, Bad, Bbc6

Target/Specificity

This mouse Bad Antibody is generated from mice immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding Ser155 of mouse Bad.

Dilution

WB~~1:500~1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Phospho-mBad(Ser155) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Phospho-mBad(Ser155) Antibody - Protein Information

Name Bad

Synonyms Bbc6

Function Promotes cell death. Successfully competes for the binding to Bcl-X(L), Bcl-2 and Bcl-W,

thereby affecting the level of heterodimerization of these proteins with BAX. Can reverse the death repressor activity of Bcl-X(L), but not that of Bcl-2. Appears to act as a link between growth factor receptor signaling and the apoptotic pathways.

Cellular Location


Mitochondrion outer membrane. Cytoplasm. Note=Colocalizes with HIF3A isoform 2 in the cytoplasm (PubMed:21546903). Upon phosphorylation, locates to the cytoplasm.

Phospho-mBad(Ser155) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Phospho-mBad(Ser155) Antibody - Images



Western analysis of cell extracts from 293 cells transfected with Flag-Bad, using Phospho-mBad-Ser155 Antibody.

Phospho-mBad(Ser155) Antibody - References

Suppression of B-cell lymphomagenesis by the BH3-only proteins Bmf and Bad. Frenzel A, et al. Blood, 2010 Feb 4. PMID 19965635.

GLP-1 mediates antiapoptotic effect by phosphorylating Bad through a beta-arrestin 1-mediated ERK1/2 activation in pancreatic beta-cells. Quoyer J, et al. J Biol Chem, 2010 Jan 15. PMID 19915011.

Identification of novel in vivo phosphorylation sites of the human proapoptotic protein BAD: pore-forming activity of BAD is regulated by phosphorylation. Polzien L, et al. J Biol Chem, 2009 Oct 9. PMID 19667065.

Perinatal survivin is essential for the establishment of pancreatic beta cell mass in mice. Wu X, et al. Diabetologia, 2009 Oct. PMID 19644667.

MEK/ERK-mediated phosphorylation of Bim is required to ensure survival of T and B lymphocytes

during mitogenic stimulation. O'Reilly LA, et al. J Immunol, 2009 Jul 1. PMID 19542438.