

VEGFC Antibody
Mouse Monoclonal Antibody (Mab)
Catalog # AM1886B

Specification

VEGFC Antibody - Product Information

Application	WB,E
Primary Accession	P49767
Other Accession	NP_005420.1
Reactivity	Human, Mouse, Rat
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1,K
Calculated MW	46883

VEGFC Antibody - Additional Information

Gene ID 7424

Other Names

Vascular endothelial growth factor C, VEGF-C, Flt4 ligand, Flt4-L, Vascular endothelial growth factor-related protein, VRP, VEGFC

Target/Specificity

This VEGFC monoclonal antibody is generated from mouse immunized with VEGFC recombinant protein.

Dilution

WB~~1:2000

Format

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

VEGFC Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

VEGFC Antibody - Protein Information

Name VEGFC

Function Growth factor active in angiogenesis, and endothelial cell growth, stimulating their proliferation and migration and also has effects on the permeability of blood vessels. May function in angiogenesis of the venous and lymphatic vascular systems during embryogenesis, and also in

the maintenance of differentiated lymphatic endothelium in adults. Binds and activates KDR/VEGFR2 and FLT4/VEGFR3 receptors.

Cellular Location

Secreted.

Tissue Location


Expressed in the spleen (PubMed:8700872, PubMed:9247316). Expressed in the lymph node, thymus, appendix and bone marrow (PubMed:9247316). Expressed in the heart, placenta, skeletal muscle, ovary and small intestine (PubMed:8617204, PubMed:8700872) Expressed in the prostate, testis and colon (PubMed:8700872)

VEGFC Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

VEGFC Antibody - Images



Western blot analysis of lysates from 293, CEM cell line and mouse heart, rat heart tissue lysates (from left to right), using VEGFC Antibody (Cat. # AM1886b). AM1886b was diluted at 1:1000 at each lane. A goat anti-mouse IgG H&L(HRP) at 1:3000 dilution was used as the secondary antibody. Lysates at 35µg per lane.



All lanes : Anti-VEGFC at 1:2000 dilution Lane 1: CCRF-CEM whole cell lysate Lane 2: 293 whole cell lysate Lane 3: HepG2 whole cell lysate Lane 4: mouse heart lysate Lane 5: rat heart lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 47 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

VEGFC Antibody - Background

The protein encoded by this gene is a member of the platelet-derived growth factor/vascular endothelial growth factor (PDGF/VEGF) family, is active in angiogenesis and endothelial cell growth, and can also affect the permeability of blood vessels. This secreted protein undergoes a complex proteolytic maturation, generating multiple processed forms which bind and activate VEGFR-3 receptors. Only the fully processed form can bind and activate VEGFR-2 receptors. This protein is structurally and functionally similar to vascular endothelial growth factor D. [provided by RefSeq].

VEGFC Antibody - References

Chen, X., et al. *Cancer Sci.* 101(11):2384-2390(2010)
Romero, R., et al. *Am. J. Obstet. Gynecol.* 203 (4), 361 (2010) :
Bailey, S.D., et al. *Diabetes Care* 33(10):2250-2253(2010)
Deguchi, K., et al. *Anticancer Res.* 30(6):2361-2366(2010)
Johnatty, S.E., et al. *PLoS Genet.* 6 (7), E1001016 (2010) :