

AKT1S1 Antibody
Mouse Monoclonal Antibody (Mab)
Catalog # AM2018b

Specification

AKT1S1 Antibody - Product Information

| | |
|-------------------|--------------------------------|
| Application | WB,E |
| Primary Accession | O96B36 |
| Other Accession | NP_001092102.1 |
| Reactivity | Human |
| Host | Mouse |
| Clonality | Monoclonal |
| Isotype | IgG1 |

AKT1S1 Antibody - Additional Information

Gene ID 84335

Other Names

Proline-rich AKT1 substrate 1, 40 kDa proline-rich AKT substrate, AKT1S1
{ECO:0000312|EMBL:AAH160431}

Target/Specificity

Purified His-tagged AKT1S1 protein(Fragment) was used to produced this monoclonal antibody.

Dilution

WB~~1:100

Format

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

AKT1S1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

AKT1S1 Antibody - Protein Information

Name AKT1S1 {ECO:0000312|EMBL:AAH16043.1}

Function Negative regulator of the mechanistic target of rapamycin complex 1 (mTORC1), an evolutionarily conserved central nutrient sensor that stimulates anabolic reactions and macromolecule biosynthesis to promote cellular biomass generation and growth (PubMed:[17277771](#), PubMed:[17386266](#), PubMed:[17510057](#), PubMed:[29236692](#)). In absence of insulin and nutrients, AKT1S1 associates with the mTORC1 complex and directly inhibits mTORC1

activity by blocking the MTOR substrate- recruitment site (PubMed:[29236692](#)). In response to insulin and nutrients, AKT1S1 dissociates from mTORC1 (PubMed:[17386266](#), PubMed:[18372248](#)). Its activity is dependent on its phosphorylation state and binding to 14-3-3 (PubMed:[16174443](#), PubMed:[18372248](#)). May also play a role in nerve growth factor-mediated neuroprotection (By similarity).

Cellular Location

Cytoplasm, cytosol {ECO:0000250|UniProtKB:Q9D1F4}. Note=Found in the cytosolic fraction of the brain. {ECO:0000250|UniProtKB:Q9D1F4}

Tissue Location


Widely expressed with highest levels of expression in liver and heart. Expressed at higher levels in cancer cell lines (e.g. A-549 and HeLa) than in normal cell lines (e.g. HEK293)

AKT1S1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

AKT1S1 Antibody - Images



AKT1S1 Antibody (Cat. #AM2018b) western blot analysis in A375 cell line lysates (35µg/lane). This demonstrates the AKT1S1 antibody detected the AKT1S1 protein (arrow).

AKT1S1 Antibody - Background

AKT1S1 is a proline-rich substrate of AKT (MIM 164730) that binds 14-3-3 protein (see YWHAH, MIM 113508) when phosphorylated (Kovacina et al., 2003 [PubMed 12524439]). [supplied by OMIM].

AKT1S1 Antibody - References

Wang, L., et al. J. Biol. Chem. 283(23):15619-15627(2008)
Fonseca, B.D., et al. Biochem. J. 411(1):141-149(2008)
Fonseca, B.D., et al. J. Biol. Chem. 282(34):24514-24524(2007)
Wang, L., et al. J. Biol. Chem. 282(27):20036-20044(2007)
Sancak, Y., et al. Mol. Cell 25(6):903-915(2007)