

CRP Antibody
Mouse Monoclonal Antibody (Mab)
Catalog # AM2041b

Specification

CRP Antibody - Product Information

Application	WB,E
Primary Accession	P02741
Other Accession	NP_000558.2
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG3
Calculated MW	25039
Antigen Region	8-35

CRP Antibody - Additional Information

Gene ID 1401

Other Names

C-reactive protein, C-reactive protein(1-205), CRP, PTX1

Target/Specificity

This CRP antibody is generated from mice immunized with a KLH conjugated synthetic peptide between 8-35 amino acids from human CRP.

Dilution

WB~~1:500~1000

Format

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Euglobin precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CRP Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

CRP Antibody - Protein Information

Name CRP

Synonyms PTX1

Function Displays several functions associated with host defense: it promotes agglutination,

bacterial capsular swelling, phagocytosis and complement fixation through its calcium-dependent binding to phosphorylcholine. Can interact with DNA and histones and may scavenge nuclear material released from damaged circulating cells.

Cellular Location
Secreted.


Tissue Location
Found in plasma.

CRP Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CRP Antibody - Images



CRP Antibody (Cat. #AM2041b) western blot analysis in NCI-H460 cell line lysates (35µg/lane). This demonstrates the CRP antibody detected the CRP protein (arrow).

CRP Antibody - Background

The protein encoded by this gene belongs to the pentaxin family. It is involved in several host defense related functions based on its ability to recognize foreign pathogens and damaged cells of the host and to initiate their elimination by interacting with humoral and cellular effector systems in the blood. Consequently, the level of this protein in plasma increases greatly during acute phase response to tissue injury, infection, or other inflammatory stimuli.

CRP Antibody - References

Brindle, E., et al. J. Immunol. Methods 362 (1-2), 112-120 (2010) :
Lim, L.S., et al. Invest. Ophthalmol. Vis. Sci. 51(9):4458-4463(2010)
Marott, S.C., et al. J. Am. Coll. Cardiol. 56(10):789-795(2010)
Szczesny, G., et al. Arch Orthop Trauma Surg (2010) In press :
Kuhlenbaeumer, G., et al. Cerebrovasc. Dis. 30(5):476-482(2010)