

IGHG1 Antibody (Ascites)
Mouse Monoclonal Antibody (Mab)
Catalog # AM2055a

Specification

IGHG1 Antibody (Ascites) - Product Information

Application	WB,E
Primary Accession	P01857
Other Accession	P01860
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	43912
Antigen Region	154-180

IGHG1 Antibody (Ascites) - Additional Information

Other Names

Ig gamma-1 chain C region, IGHG1

Target/Specificity

This IGHG1 antibody is generated from mice immunized with a KLH conjugated synthetic peptide between 154-180 amino acids from human IGHG1.

Dilution

WB~~1:100~1600

Format

Mouse monoclonal antibody supplied in crude ascites with 0.09% (W/V) sodium azide.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

IGHG1 Antibody (Ascites) is for research use only and not for use in diagnostic or therapeutic procedures.

IGHG1 Antibody (Ascites) - Protein Information

Name IGHG1 {ECO:0000303|PubMed:11340299, ECO:0000303|Ref.13}

Function Constant region of immunoglobulin heavy chains. Immunoglobulins, also known as antibodies, are membrane-bound or secreted glycoproteins produced by B lymphocytes. In the recognition phase of humoral immunity, the membrane-bound immunoglobulins serve as receptors which, upon binding of a specific antigen, trigger the clonal expansion and differentiation of B lymphocytes into immunoglobulins- secreting plasma cells. Secreted

immunoglobulins mediate the effector phase of humoral immunity, which results in the elimination of bound antigens (PubMed:[20176268](#), PubMed:[22158414](#)). The antigen binding site is formed by the variable domain of one heavy chain, together with that of its associated light chain. Thus, each immunoglobulin has two antigen binding sites with remarkable affinity for a particular antigen. The variable domains are assembled by a process called V-(D)-J rearrangement and can then be subjected to somatic hypermutations which, after exposure to antigen and selection, allow affinity maturation for a particular antigen (PubMed:[17576170](#), PubMed:[20176268](#)). Mediates IgG effector functions on monocytes triggering ADCC of virus- infected cells.

Cellular Location


[Isoform 1]: Secreted

IGHG1 Antibody (Ascites) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

IGHG1 Antibody (Ascites) - Images



IGHG1 Antibody (Cat. #AM2055a) western blot analysis in HepG2 cell line lysates (35µg/lane). This demonstrates the IGHG1 antibody detected the IGHG1 protein (arrow).

IGHG1 Antibody (Ascites) - Background

The function of this protein remains unknown.