

HGF Antibody
Mouse Monoclonal Antibody (Mab)
Catalog # AM2064b

Specification

HGF Antibody - Product Information

Application	WB,E
Primary Accession	P14210
Other Accession	NP_000592.3
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	83134
Antigen Region	521-554

HGF Antibody - Additional Information

Gene ID 3082

Other Names

Hepatocyte growth factor, Hepatopoietin-A, Scatter factor, SF, Hepatocyte growth factor alpha chain, Hepatocyte growth factor beta chain, HGF, HPTA

Target/Specificity

This HGF antibody is generated from mice immunized with a KLH conjugated synthetic peptide between 521-554 amino acids from human HGF.

Dilution

WB~~1:4000

Format

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

HGF Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

HGF Antibody - Protein Information

Name HGF

Synonyms HPTA


Function Potent mitogen for mature parenchymal hepatocyte cells, seems to be a hepatotrophic factor, and acts as a growth factor for a broad spectrum of tissues and cell types (PubMed:[20624990](#)). Activating ligand for the receptor tyrosine kinase MET by binding to it and promoting its dimerization (PubMed:[15167892](#), PubMed:[20977675](#)). Activates MAPK signaling following Tmprss13 cleavage and activation (PubMed:[20977675](#)).

HGF Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

HGF Antibody - Images



All lanes : Anti-Human HGF C-term at 1:1000 dilution Lane 1: HepG2 whole cell lysate Lane 2: Ramos whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 83 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Anti-Human HGF C-term at 1:4000 dilution + Human brain lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 83 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

HGF Antibody - Background

Hepatocyte growth factor regulates cell growth, cell motility, and morphogenesis by activating a tyrosine kinase signaling cascade after binding to the proto-oncogenic c-Met receptor. Hepatocyte growth factor is secreted by mesenchymal cells and acts as a multi-functional cytokine on cells of mainly epithelial origin. Its ability to stimulate mitogenesis, cell motility, and matrix invasion gives it a central role in angiogenesis, tumorigenesis, and tissue regeneration. It is secreted as a single inactive polypeptide and is cleaved by serine proteases into a 69-kDa alpha-chain and 34-kDa beta-chain. A disulfide bond between the alpha and beta chains produces the active, heterodimeric molecule. The protein belongs to the plasminogen subfamily of S1 peptidases but has no detectable protease activity. Alternative splicing of this gene produces multiple transcript variants encoding different isoforms. [provided by RefSeq].

HGF Antibody - References

- Shi, E., et al. *Anesthesiology* 113(5):1109-1117(2010)
- Hoot, K.E., et al. *J. Clin. Invest.* 120(10):3606-3616(2010)
- Bailey, S.D., et al. *Diabetes Care* 33(10):2250-2253(2010)
- Togawa, A., et al. *Biochem. Biophys. Res. Commun.* 400(2):271-277(2010)
- Li, F., et al. *J. Biol. Chem.* 285(36):27673-27685(2010)