

**CYP1A2 Antibody (Ascites)**  
**Mouse Monoclonal Antibody (Mab)**  
**Catalog # AM2066a**

**Specification**

---

**CYP1A2 Antibody (Ascites) - Product Information**

Application	WB,E
Primary Accession	<a href="#">P05177</a>
Other Accession	<a href="#">NP_000752.2</a>
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG3
Calculated MW	58407
Antigen Region	255-282

**CYP1A2 Antibody (Ascites) - Additional Information**

**Gene ID** 1544

**Other Names**

Cytochrome P450 1A2, CYP1A2, Cytochrome P(3)450, Cytochrome P450 4, Cytochrome P450-P3, CYP1A2

**Target/Specificity**

This CYP1A2 antibody is generated from mice immunized with a KLH conjugated synthetic peptide between 255-282 amino acids from human CYP1A2.

**Dilution**

WB~~1:500~2000

**Format**

Mouse monoclonal antibody supplied in crude ascites with 0.09% (W/V) sodium azide.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

CYP1A2 Antibody (Ascites) is for research use only and not for use in diagnostic or therapeutic procedures.

**CYP1A2 Antibody (Ascites) - Protein Information**

**Name** CYP1A2 {ECO:0000303|PubMed:2575218, ECO:0000312|HGNC:HGNC:2596}

**Function** A cytochrome P450 monooxygenase involved in the metabolism of various endogenous substrates, including fatty acids, steroid hormones and vitamins (PubMed:[10681376](#),

PubMed:[11555828](#), PubMed:[12865317](#), PubMed:[19965576](#), PubMed:[9435160](#)). Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (NADPH--hemoprotein reductase) (PubMed:[10681376](#), PubMed:[11555828](#), PubMed:[12865317](#), PubMed:[19965576](#), PubMed:[9435160](#)). Catalyzes the hydroxylation of carbon-hydrogen bonds (PubMed:[11555828](#), PubMed:[12865317](#)). Exhibits high catalytic activity for the formation of hydroxysteroids from estrone (E1) and 17beta- estradiol (E2), namely 2-hydroxy E1 and E2 (PubMed:[11555828](#), PubMed:[12865317](#)). Metabolizes cholesterol toward 25-hydroxycholesterol, a physiological regulator of cellular cholesterol homeostasis (PubMed:[21576599](#)). May act as a major enzyme for all-trans retinoic acid biosynthesis in the liver. Catalyzes two successive oxidative transformation of all-trans retinol to all-trans retinal and then to the active form all-trans retinoic acid (PubMed:[10681376](#)). Primarily catalyzes stereoselective epoxidation of the last double bond of polyunsaturated fatty acids (PUFA), displaying a strong preference for the (R,S) stereoisomer (PubMed:[19965576](#)). Catalyzes bisallylic hydroxylation and omega-1 hydroxylation of PUFA (PubMed:[9435160](#)). May also participate in eicosanoids metabolism by converting hydroperoxide species into oxo metabolites (lipoxygenase-like reaction, NADPH- independent) (PubMed:[21068195](#)). Plays a role in the oxidative metabolism of xenobiotics. Catalyzes the N-hydroxylation of heterocyclic amines and the O-deethylation of phenacetin (PubMed:[14725854](#)). Metabolizes caffeine via N3-demethylation (Probable).

#### Cellular Location

Endoplasmic reticulum membrane; Peripheral membrane protein. Microsome membrane; Peripheral membrane protein

#### Tissue Location


Liver.


### CYP1A2 Antibody (Ascites) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### CYP1A2 Antibody (Ascites) - Images





CYP1A2 Antibody (Cat. #AM2066a) western blot analysis in SK-BR-3 cell line lysates (35µg/lane). This demonstrates the CYP1A2 antibody detected the CYP1A2 protein (arrow).

### **CYP1A2 Antibody (Ascites) - Background**

This gene encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. The protein encoded by this gene localizes to the endoplasmic reticulum and its expression is induced by some polycyclic aromatic hydrocarbons (PAHs), some of which are found in cigarette smoke. The enzyme's endogenous substrate is unknown; however, it is able to metabolize some PAHs to carcinogenic intermediates. Other xenobiotic substrates for this enzyme include caffeine, aflatoxin B1, and acetaminophen. The transcript from this gene contains four Alu sequences flanked by direct repeats in the 3' untranslated region.

### **CYP1A2 Antibody (Ascites) - References**

Gentile, G., et al. *J Headache Pain* 11(5):431-435(2010)  
Uslu, A., et al. *BMB Rep* 43(8):530-534(2010)  
Wang, X., et al. *J. Pharm. Pharmacol.* 62(8):1077-1083(2010)  
Schmidt, R.J., et al. *Birth Defects Res. Part A Clin. Mol. Teratol.* 88(7):560-569(2010)  
Jiang, Z., et al. *Pharmacogenet. Genomics* 16(5):359-367(2006)