

STRA8 Antibody
Mouse Monoclonal Antibody (Mab)
Catalog # AM2091a

Specification

STRA8 Antibody - Product Information

Application	WB,E
Primary Accession	O7Z7C7
Other Accession	NP_872295.1
Reactivity	Human, Mouse
Host	Mouse
Clonality	Monoclonal
Isotype	IgM
Antigen Region	31-59

STRA8 Antibody - Additional Information

Gene ID 346673

Other Names

Stimulated by retinoic acid gene 8 protein homolog, STRA8 {ECO:0000312|EMBL:AAP471631}

Target/Specificity

This STRA8 antibody is generated from mice immunized with a KLH conjugated synthetic peptide between 31-59 amino acids from human STRA8.

Dilution

WB~~1:8000

Format

Mouse monoclonal antibody supplied in crude ascites with 0.09% (W/V) sodium azide.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

STRA8 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

STRA8 Antibody - Protein Information

Name STRA8 ([HGNC:30653](#))

Function Meiosis-inducer required for the transition into meiosis for both female and male germ cells. In female germ cells, acts downstream of ZGLP1 as a key effector of the meiotic program: required for premeiotic DNA replication and subsequent events in meiotic prophase. During spermatogenesis, next to its role in meiotic initiation, promotes (but is not required for) spermatogonial differentiation. In complex with MEIOSIN, directly activates the transcription of a

subset of critical meiotic genes playing a central role in cell-cycle switching from mitosis to meiosis.

Cellular Location

Cytoplasm {ECO:0000250|UniProtKB:P70278}. Nucleus {ECO:0000250|UniProtKB:P70278}.
Note=Shuttles between nucleus and cytoplasm. Nuclear export is XPO1-dependent {ECO:0000250|UniProtKB:P70278}

Tissue Location


Expressed specifically in testis and fetal ovaries.

STRA8 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

STRA8 Antibody - Images



"Anti-STRA8 Antibody at 1:2000 dilution + HeLa whole cell lysate Secondary Goat Anti-mouse IgM, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 36908 Da Blocking/Dilution buffer: 5% NFDN/TBST."



"Anti-STRA8 Antibody at 1:8000 dilution + mouse bladder lysate Secondary Goat Anti-mouse IgM, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 36908 Da Blocking/Dilution buffer: 5% NFD/MTBST."

STRA8 Antibody - Background

This gene encodes a retinoic acid-responsive protein. A homologous protein in mouse has been shown to be involved in the regulation of meiotic initiation in both spermatogenesis and oogenesis, though feature differences between the mouse and human proteins suggest that these homologs are not entirely functionally equivalent. It is thought that this gene may play a role in spermatogenesis in humans.

STRA8 Antibody - References

- Le Bouffant, R., et al. Hum. Reprod. 25(10):2579-2590(2010)
- Aston, K.I., et al. Hum. Reprod. 25(6):1383-1397(2010)
- Chen, B., et al. Asian J. Androl. 11(5):557-565(2009)
- Houmard, B., et al. Biol. Reprod. 81(2):438-443(2009)
- Stouffs, K., et al. Verh. K. Acad. Geneesk. Belg. 71(3):115-139(2009)