

**HLA-B Antibody**  
**Mouse Monoclonal Antibody (Mab)**  
**Catalog # AM2114b****Specification**

---

**HLA-B Antibody - Product Information**

|                   |   |
|-------------------|---|
| Application       | <b>WB,E</b>   |
| Primary Accession | <a href="#">P03989</a>  |
| Other Accession   | <a href="#">P30490</a> , <a href="#">P30488</a> , <a href="#">P30487</a> , <a href="#">P30486</a> , <a href="#">P30485</a> ,<br><a href="#">P30483</a> , <a href="#">P30481</a> , <a href="#">P30479</a> , <a href="#">Q04826</a> , <a href="#">P18463</a> ,<br><a href="#">P30464</a> , <a href="#">P30461</a> , <a href="#">NP_005505.2</a> |
| Reactivity        | <b>Human</b>  |
| Host              | <b>Mouse</b>  |
| Clonality         | <b>Monoclonal</b>   |
| Isotype           | <b>IgM</b>  |
| Antigen Region    | <b>62-90</b>  |

**HLA-B Antibody - Additional Information****Other Names**

HLA class I histocompatibility antigen, B-27 alpha chain, MHC class I antigen B\*27, HLA-B, HLAB

**Target/Specificity**

This HLA-B antibody is generated from mice immunized with a KLH conjugated synthetic peptide between 62-90 amino acids from human HLA-B.

**Dilution**

WB~~1:500~1000

**Format**

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Euglobin precipitation followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**


HLA-B Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**HLA-B Antibody - Protein Information****HLA-B Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### HLA-B Antibody - Images



HLA-B Antibody (Cat. #AM2114b) western blot analysis in Ramos cell line lysates (35 $\mu$ g/lane). This demonstrates the HLA-B antibody detected the HLA-B protein (arrow).

### HLA-B Antibody - Background

HLA-B belongs to the HLA class I heavy chain paralogues. This class I molecule is a heterodimer consisting of a heavy chain and a light chain (beta-2 microglobulin). The heavy chain is anchored in the membrane. Class I molecules play a central role in the immune system by presenting peptides derived from the endoplasmic reticulum lumen. They are expressed in nearly all cells. The heavy chain is approximately 45 kDa and its gene contains 8 exons. Exon 1 encodes the leader peptide, exon 2 and 3 encode the alpha1 and alpha2 domains, which both bind the peptide, exon 4 encodes the alpha3 domain, exon 5 encodes the transmembrane region and exons 6 and 7 encode the cytoplasmic tail. Polymorphisms within exon 2 and exon 3 are responsible for the peptide binding specificity of each class one molecule. Typing for these polymorphisms is routinely done for bone marrow and kidney transplantation. Hundreds of HLA-B alleles have been described.

### HLA-B Antibody - References

Parsons, M.S., et al. J. Infect. Dis. 202 SUPPL 3, S356-S360 (2010) :  
Brockman, M.A., et al. J. Virol. 84(22):11937-11949(2010)  
Noble, J.A., et al. Diabetes 59(11):2972-2979(2010)  
Lipponen, K., et al. Diabetes (2010) In press :  
Liao, Y.Q., et al. Zhongguo Shi Yan Xue Ye Xue Za Zhi 18(4):1055-1058(2010)