

**H2AFX Antibody (C-term)**  
**Mouse Monoclonal Antibody (Mab)**  
**Catalog # AM2199B**

**Specification**

---

**H2AFX Antibody (C-term) - Product Information**

Application	<b>WB, IHC, IHC-P,E</b>
Primary Accession	<a href="#">P16104</a>
Reactivity	<b>Human</b>
Host	<b>Mouse</b>
Clonality	<b>Monoclonal</b>
Isotype	<b>IgG1</b>
Antigen Region	<b>115-143</b>

**H2AFX Antibody (C-term) - Additional Information**

**Gene ID** 3014

**Other Names**

Histone H2AX, H2a/x, Histone H2AX, H2AFX, H2AX

**Target/Specificity**

This H2AFX antibody is generated from mice immunized with a KLH conjugated synthetic peptide between 115-143 amino acids from the C-terminal region of human H2AFX.

**Dilution**

WB~~1:1000

IHC~~1:50

IHC-P~~1:25

**Format**

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

H2AFX Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**H2AFX Antibody (C-term) - Protein Information**

**Name** H2AX ([HGNC:4739](#))

**Function** Variant histone H2A which replaces conventional H2A in a subset of nucleosomes. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular

machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post- translational modifications of histones, also called histone code, and nucleosome remodeling. Required for checkpoint-mediated arrest of cell cycle progression in response to low doses of ionizing radiation and for efficient repair of DNA double strand breaks (DSBs) specifically when modified by C-terminal phosphorylation.

#### Cellular Location


Nucleus. Chromosome

#### H2AFX Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


#### H2AFX Antibody (C-term) - Images




Western blot analysis of lysates from 293, CEM, HepG2, Jurkat, HeLa, Raji cell line(from left to right). using H2AFX Antibody (C-term)(Cat. #AM2199b). AM2199b was diluted at 1:2000 at each lane. A goat anti-mouse IgG H&L(HRP) at 1:3000 dilution was used as the secondary antibody. Lysates at 35µg per lane.



Immunohistochemical analysis of paraffin-embedded Human Thymus section using Pink1(Cat#am2199b). am2199b was diluted at 1:50 dilution. A undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB staining.



Immunohistochemical analysis of paraffin-embedded H. prostate section using H2AFX Antibody (C-term)(Cat#AM2199b). AM2199b was diluted at 1:25 dilution. A undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB staining.



Immunohistochemical analysis of paraffin-embedded H. thymus section using H2AFX Antibody (C-term)(Cat#AM2199b). AM2199b was diluted at 1:25 dilution. A undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB staining.

### **H2AFX Antibody (C-term) - Background**

Variant histone H2A which replaces conventional H2A in a subset of nucleosomes. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling. Required for checkpoint-mediated arrest of cell cycle progression in response to low doses of ionizing radiation and for efficient repair of DNA double strand breaks (DSBs) specifically when modified by C-terminal phosphorylation.

### **H2AFX Antibody (C-term) - References**

- Stewart G.S., et al. Nature 421:961-966(2003).
- Park E.-J., et al. Nucleic Acids Res. 31:6819-6827(2003).
- Stiff T., et al. Cancer Res. 64:2390-2396(2004).
- Lukas C., et al. EMBO J. 23:2674-2683(2004).
- Kurz E.U., et al. J. Biol. Chem. 279:53272-53281(2004).