

**USP11 Antibody (C-term R565)**  
**Purified Mouse Monoclonal Antibody (Mab)**  
**Catalog # AM2260b**

**Specification**

---

**USP11 Antibody (C-term R565) - Product Information**

Application	WB, FC,E
Primary Accession	<a href="#">P51784</a>
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1, $\kappa$

**USP11 Antibody (C-term R565) - Additional Information**

**Other Names**

Ubiquitin carboxyl-terminal hydrolase 11, Deubiquitinating enzyme 11, Ubiquitin thioesterase 11, Ubiquitin-specific-processing protease 11, USP11, UHX1

**Target/Specificity**

This USP11 antibody is generated from a mouse immunized with a KLH conjugated synthetic peptide between 32-300 amino acids from the N-terminal region of human USP11.

**Dilution**

WB~~1:1000  
FC~~1:25

**Format**

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

USP11 Antibody (C-term R565) is for research use only and not for use in diagnostic or therapeutic procedures.

**USP11 Antibody (C-term R565) - Protein Information**

**Name** USP11

**Synonyms** UHX1

**Function** Protease that can remove conjugated ubiquitin from target proteins and polyubiquitin chains (PubMed:[12084015](#), PubMed:[15314155](#), PubMed:[17897950](#), PubMed:[19874889](#), PubMed:[20233726](#), PubMed:[24724799](#), PubMed:[28992046](#)). Inhibits the degradation of target

proteins by the proteasome (PubMed:[12084015](#)). Cleaves preferentially 'Lys-6' and 'Lys- 63'-linked ubiquitin chains. Has lower activity with 'Lys-11' and 'Lys- 33'-linked ubiquitin chains, and extremely low activity with 'Lys-27', 'Lys-29' and 'Lys-48'-linked ubiquitin chains (in vitro) (PubMed:[24724799](#)). Plays a role in the regulation of pathways leading to NF-kappa-B activation (PubMed:[17897950](#), PubMed:[19874889](#)). Plays a role in the regulation of DNA repair after double-stranded DNA breaks (PubMed:[15314155](#), PubMed:[20233726](#)). Acts as a chromatin regulator via its association with the Polycomb group (PcG) multiprotein PRC1-like complex; may act by deubiquitinating components of the PRC1-like complex (PubMed:[20601937](#)). Promotes cell proliferation by deubiquitinating phosphorylated E2F1 (PubMed:[28992046](#)).

#### Cellular Location


Nucleus. Cytoplasm. Chromosome. Note=Predominantly nuclear (PubMed:12084015, PubMed:15314155). Associates with chromatin (PubMed:20233726, PubMed:20601937).

#### USP11 Antibody (C-term R565) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### USP11 Antibody (C-term R565) - Images



Western blot analysis of lysates from Jurkat, HeLa, LNCaP cell line (from left to right) using USP11 Antibody (N-term) (Cat. # AM2260b). AM2260b was diluted at 1:1000 at each lane. A goat anti-mouse IgG H&L(HRP) at 1:3000 dilution was used as the secondary antibody. Lysates at 35µg per lane.



Flow cytometric analysis of Hela cells using USP11 Antibody (C-term R565)(green, Cat#AM2260b) compared to an isotype control of mouse IgG1(blue). AM2260b was diluted at 1:25 dilution. An Alexa Fluor® 488 goat anti-mouse IgG at 1:400 dilution was used as the secondary antibody.

#### **USP11 Antibody (C-term R565) - Background**

Protease that can remove conjugated ubiquitin from target proteins and polyubiquitin chains. Inhibits the degradation of target proteins by the proteasome. Plays a role in the regulation of pathways leading to NF-kappa-B activation. Plays a role in the regulation of DNA repair after double-stranded DNA breaks.

#### **USP11 Antibody (C-term R565) - References**

Ross M.T.,et al.Nature 434:325-337(2005).  
Ideguchi H.,et al.Biochem. J. 367:87-95(2002).  
Swanson D.A.,et al.Hum. Mol. Genet. 5:533-538(1996).  
Schoenfeld A.R.,et al.Mol. Cell. Biol. 24:7444-7455(2004).  
Yamaguchi T.,et al.J. Biol. Chem. 282:33943-33948(2007).