

MYB Antibody (Center)
Purified Mouse Monoclonal Antibody (Mab)
Catalog # AM2268b

Specification

MYB Antibody (Center) - Product Information

| | |
|-------------------|---------------------------------|
| Application | WB,E |
| Primary Accession | P10242 |
| Reactivity | Human |
| Host | Mouse |
| Clonality | Monoclonal |
| Isotype | IgG1,κ |

MYB Antibody (Center) - Additional Information

Gene ID 4602

Other Names

Transcriptional activator Myb, Proto-oncogene c-Myb, MYB

Target/Specificity

This MYB antibody is generated from a mouse immunized with a KLH conjugated synthetic peptide between 1-185 amino acids from the Central region of human MYB.

Dilution

WB~~1:2000

Format

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

MYB Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

MYB Antibody (Center) - Protein Information

Name MYB

Function Transcriptional activator; DNA-binding protein that specifically recognize the sequence 5'-YAAC[GT]G-3'. Plays an important role in the control of proliferation and differentiation of hematopoietic progenitor cells.

Cellular Location


Nucleus {ECO:0000255|PROSITE-ProRule:PRU00625, ECO:0000269|PubMed:19646965}

MYB Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MYB Antibody (Center) - Images



All lanes : Anti-MYB Antibody (Center) at 1:2000 dilution Lane 1: Jurkat whole cell lysate Lane 2: CCRF-CEM whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 80 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

MYB Antibody (Center) - Background

Transcriptional activator; DNA-binding protein that specifically recognize the sequence 5'-YAAC[GT]G-3'. Plays an important role in the control of proliferation and differentiation of hematopoietic progenitor cells.

MYB Antibody (Center) - References

- Majello B.,et al.Proc. Natl. Acad. Sci. U.S.A. 83:9636-9640(1986).
Westin E.H.,et al.Oncogene 5:1117-1124(1990).
Close J.P.,et al.BMC Genomics 5:33-33(2004).
Westin E.H.,et al.Submitted (MAR-1995) to the EMBL/GenBank/DDBJ databases.
Gaillard C.,et al.Submitted (NOV-1998) to the EMBL/GenBank/DDBJ databases.