

CRYAB Antibody
Purified Mouse Monoclonal Antibody (Mab)
Catalog # AM8424b

Specification

CRYAB Antibody - Product Information

Application	WB,E
Primary Accession	P02511
Reactivity	Human, Mouse, Rat
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1, κ
Calculated MW	20159

CRYAB Antibody - Additional Information

Gene ID 1410

Other Names

Alpha-crystallin B chain, Alpha(B)-crystallin, Heat shock protein beta-5, HspB5, Renal carcinoma antigen NY-REN-27, Rosenthal fiber component, CRYAB, CRYA2

Target/Specificity

This CRYAB antibody is generated from a mouse immunized with a recombination protein from the human region of human CRYAB.

Dilution

WB~~1:1000

Format

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CRYAB Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

CRYAB Antibody - Protein Information

Name CRYAB ([HGNC:2389](#))

Synonyms CRYA2, HSPB5

Function May contribute to the transparency and refractive index of the lens. Has chaperone-like activity, preventing aggregation of various proteins under a wide range of stress conditions. In lens

epithelial cells, stabilizes the ATP6V1A protein, preventing its degradation by the proteasome (By similarity).

Cellular Location

Cytoplasm. Nucleus Secreted. Lysosome {ECO:0000250|UniProtKB:P23927}. Note=Translocates to the nucleus during heat shock and resides in sub-nuclear structures known as SC35 speckles or nuclear splicing speckles (PubMed:19464326). Localizes at the Z- bands and the intercalated disk in cardiomyocytes (PubMed:28493373) Can be secreted; the secretion is dependent on protein unfolding and facilitated by the cargo receptor TMED10; it results in protein translocation from the cytoplasm into the ERGIC (endoplasmic reticulum- Golgi intermediate compartment) followed by vesicle entry and secretion (PubMed:32272059).

Tissue Location


Lens as well as other tissues (PubMed:2387586, PubMed:838078). Expressed in myocardial tissue (PubMed:28493373)

CRYAB Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CRYAB Antibody - Images



Western blot analysis of lysates from U-87 MG, MCF-7 cell line, mouse heart and skeletal muscle, rat skeletal muscle tissue lysates (from left to right), using CRYAB Antibody (Cat. #AM8424b). AM8424b was diluted at 1:1000 at each lane. A goat anti-mouse IgG H&L (HRP) at 1:3000 dilution was used as the secondary antibody. Lysates at 35µg per lane.

CRYAB Antibody - Background

May contribute to the transparency and refractive index of the lens. Has chaperone-like activity, preventing aggregation of various proteins under a wide range of stress conditions.

CRYAB Antibody - References

- Kramps J.A., et al. FEBS Lett. 74:82-84(1977).
Dubin R.A., et al. Genomics 7:594-601(1990).
Iwaki A., et al. Neurosci. Lett. 140:89-92(1992).
Yu W., et al. Submitted (JUN-1997) to the EMBL/GenBank/DDBJ databases.
Ota T., et al. Nat. Genet. 36:40-45(2004).