

JUN antibody
Purified Mouse Monoclonal Antibody (Mab)
Catalog # AM8428b

Specification

JUN antibody - Product Information

| | |
|-------------------|------------------------|
| Application | WB,E |
| Primary Accession | P05412 |
| Reactivity | Human, Mouse |
| Host | Mouse |
| Clonality | Monoclonal |
| Isotype | IgG1,k |
| Calculated MW | 35676 |

JUN antibody - Additional Information

Gene ID 3725

Other Names

Transcription factor AP-1, Activator protein 1, AP1, Proto-oncogene c-Jun, V-jun avian sarcoma virus 17 oncogene homolog, p39, JUN

Target/Specificity

This antibody is generated from a mouse immunized with a KLH conjugated synthetic peptide between amino acids from human.

Dilution

WB~~1:1000

Format

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

JUN antibody is for research use only and not for use in diagnostic or therapeutic procedures.

JUN antibody - Protein Information

Name JUN

Function Transcription factor that recognizes and binds to the AP-1 consensus motif 5'-TGA[GC]TCA-3' (PubMed:[10995748](#), PubMed:[22083952](#)). Heterodimerizes with proteins of the FOS family to form an AP-1 transcription complex, thereby enhancing its DNA binding activity to the AP-1 consensus sequence 5'-TGA[GC]TCA-3' and enhancing its transcriptional activity (By

similarity). Together with FOSB, plays a role in activation-induced cell death of T cells by binding to the AP-1 promoter site of FASLG/CD95L, and inducing its transcription in response to activation of the TCR/CD3 signaling pathway (PubMed:[12618758](#)). Promotes activity of NR5A1 when phosphorylated by HIPK3 leading to increased steroidogenic gene expression upon cAMP signaling pathway stimulation (PubMed:[17210646](#)). Involved in activated KRAS-mediated transcriptional activation of USP28 in colorectal cancer (CRC) cells (PubMed:[24623306](#)). Binds to the USP28 promoter in colorectal cancer (CRC) cells (PubMed:[24623306](#)).

Cellular Location

Nucleus.

Tissue Location


Expressed in the developing and adult prostate and prostate cancer cells.

JUN antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

JUN antibody - Images



Western blot analysis of lysates from 293, mouse NIH/3T3 cell line (from left to right), using JUN Antibody (Cat. #AM8428b). AM8428b was diluted at 1:1000 at each lane. A goat anti-mouse IgG H&L(HRP) at 1:3000 dilution was used as the secondary antibody. Lysates at 35µg per lane.

JUN antibody - Background

Transcription factor that recognizes and binds to the enhancer heptamer motif 5'-TGA[CG]TCA-3'. Promotes activity of NR5A1 when phosphorylated by HIPK3 leading to increased steroidogenic gene

expression upon cAMP signaling pathway stimulation.

JUN antibody - References

- Hattori K., et al. Proc. Natl. Acad. Sci. U.S.A. 85:9148-9152(1988).
Bohmann D., et al. Science 238:1386-1392(1987).
Kalnine N., et al. Submitted (OCT-2004) to the EMBL/GenBank/DDBJ databases.
Gregory S.G., et al. Nature 441:315-321(2006).
Boyle W.J., et al. Cell 64:573-584(1991).