

VWF Antibody
Purified Mouse Monoclonal Antibody (Mab)
Catalog # AM8430b

Specification

VWF Antibody - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	P04275
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	309265

VWF Antibody - Additional Information

Gene ID 7450

Other Names

von Willebrand factor, vWF, von Willebrand antigen 2, von Willebrand antigen II, VWF, F8VWF

Target/Specificity

This antibody is generated from a mouse immunized with .VWF recombinant protein.

Dilution

WB~~1□2000

IHC-P~~1:25

FC~~1:100

Format

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

VWF Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

VWF Antibody - Protein Information

Name VWF

Synonyms F8VWF

Function Important in the maintenance of hemostasis, it promotes adhesion of platelets to the sites of vascular injury by forming a molecular bridge between sub-endothelial collagen matrix and

platelet- surface receptor complex GPIb-IX-V. Also acts as a chaperone for coagulation factor VIII, delivering it to the site of injury, stabilizing its heterodimeric structure and protecting it from premature clearance from plasma.

Cellular Location

Secreted. Secreted, extracellular space, extracellular matrix. Note=Localized to storage granules

Tissue Location


Plasma.

VWF Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


VWF Antibody - Images



Anti-VWF at dilution + VWF whole cell lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 309 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.



Immunohistochemical analysis of paraffin-embedded H. kidney section using VWF(Cat#AM8430b). AM8430b was diluted at 1:25 dilution. A peroxidase-conjugated goat anti-mouse IgG at 1:400 dilution was used as the secondary antibody, followed by DAB staining.



Flow cytometric analysis of K562 cells using VWF(green, Cat#AM8430b) compared to an isotype control of mouse IgG1(blue). AP20600c was diluted at 1:100 dilution. An Alexa Fluor® 488 goat anti-mouse IgG at 1:400 dilution was used as the secondary antibody.

VWF Antibody - Background

Important in the maintenance of hemostasis, it promotes adhesion of platelets to the sites of vascular injury by forming a molecular bridge between sub-endothelial collagen matrix and platelet-surface receptor complex GPIb-IX-V. Also acts as a chaperone for coagulation factor VIII, delivering it to the site of injury, stabilizing its heterodimeric structure and protecting it from premature clearance from plasma.

VWF Antibody - References

- Bonthron D.,et al.Nucleic Acids Res. 14:7125-7128(1986).
- Mancuso D.J.,et al.J. Biol. Chem. 264:19514-19527(1989).
- Scherer S.E.,et al.Nature 440:346-351(2006).
- Verweij C.L.,et al.EMBO J. 5:1839-1847(1986).
- Verweij C.L.,et al.EMBO J. 5:3074-3074(1986).