

**LAMP1 Antibody**  
**Purified Mouse Monoclonal Antibody (Mab)**  
**Catalog # AM8465b**

**Specification**

---

**LAMP1 Antibody - Product Information**

Application	<b>WB, IF, IHC-P,E</b>
Primary Accession	<a href="#">P11279</a>
Reactivity	<b>Human, Green Monkey</b>
Host	<b>Mouse</b>
Clonality	<b>monoclonal</b>
Isotype	<b>IgG1,k</b>

**LAMP1 Antibody - Additional Information**

**Gene ID** 3916

**Other Names**

Lysosome-associated membrane glycoprotein 1, LAMP-1, Lysosome-associated membrane protein 1, CD107 antigen-like family member A, CD107a, LAMP1

**Target/Specificity**

This LAMP1 antibody is generated from a mouse immunized with a recombinant protein.

**Dilution**

WB~~1:2000

IF~~1:25

IHC-P~~1:25

**Format**

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

LAMP1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**LAMP1 Antibody - Protein Information**

**Name** LAMP1 {ECO:0000303|PubMed:23632890, ECO:0000312|HGNC:HGNC:6499}

**Function** Lysosomal membrane glycoprotein which plays an important role in lysosome biogenesis, lysosomal pH regulation, autophagy and cholesterol homeostasis (PubMed:[37390818](#)). Acts as an important regulator of lysosomal lumen pH regulation by acting as a direct inhibitor of the proton channel TMEM175, facilitating lysosomal acidification for optimal hydrolase activity

(PubMed:[37390818](#)). Also plays an important role in NK-cells cytotoxicity (PubMed:[2022921](#), PubMed:[23632890](#)). Mechanistically, participates in cytotoxic granule movement to the cell surface and perforin trafficking to the lytic granule (PubMed:[23632890](#)). In addition, protects NK-cells from degranulation-associated damage induced by their own cytotoxic granule content (PubMed:[23847195](#)). Presents carbohydrate ligands to selectins (PubMed:[7685349](#)).

#### Cellular Location


Lysosome membrane; Single-pass type I membrane protein. Endosome membrane; Single-pass type I membrane protein. Late endosome membrane; Single-pass type I membrane protein. Cell membrane; Single-pass type I membrane protein. Cytolytic granule membrane; Single-pass type I membrane protein. Note=This protein shuttles between lysosomes, endosomes, and the plasma membrane (By similarity). Colocalizes with OSBPL1A at the late endosome (PubMed:16176980). {ECO:0000250|UniProtKB:P05300, ECO:0000269|PubMed:16176980, ECO:0000269|PubMed:17897319}

#### LAMP1 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


#### LAMP1 Antibody - Images




All lanes : Anti-LAMP1 Antibody at 1:1000 dilution (upper) Lane 1: HT1080 Lane 2: HT1080-Knockout Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Mouse IgG, (H+L), Peroxidase conjugated (ASP1613) at 1/8000 dilution. Predicted band size : 90-120 kDa



Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized HeLa (human cervical epithelial adenocarcinoma cell line) cells labeling LAMP1 with AM8465b at 1/25 dilution, followed by Dylight® 488-conjugated goat anti-mouse IgG (NA166821) secondary antibody at 1/200 dilution (green). Immunofluorescence image showing cytoplasm staining on HeLa cell line. Cytoplasmic actin is detected with Dylight® 554 Phalloidin (PD18466410) at 1/100 dilution (red). The nuclear counter stain is DAPI (blue).



All lanes : Anti-LAMP1 Antibody at 1:2000 dilution Lane 1: A431 whole cell lysates Lane 2: HeLa whole cell lysates Lane 3: Jurkat whole cell lysates Lane 4: HT-1080 whole cell lysates Lane 5: COS-7 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 45 kDa Blocking/Dilution buffer: 5% NFDN/TBST.



AM8465b staining LAMP1 in human kidney sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hour at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.

#### **LAMP1 Antibody - Background**

Presents carbohydrate ligands to selectins. Also implicated in tumor cell metastasis.

#### **LAMP1 Antibody - References**

Fukuda M., et al. *J. Biol. Chem.* 263:18920-18928(1988).  
Ota T., et al. *Nat. Genet.* 36:40-45(2004).  
Dunham A., et al. *Nature* 428:522-528(2004).  
Mural R.J., et al. Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.  
Mane S.M., et al. *Arch. Biochem. Biophys.* 268:360-378(1989).

#### **LAMP1 Antibody - Citations**

- [Effect of Protein O-Mannosyltransferase \(MSMEG\\_5447\) on and Its Survival in Macrophages](#)
- [L-3-n-Butylphthalide improves synaptic and dendritic spine plasticity and ameliorates neurite pathology in Alzheimer's disease mouse model and cultured hippocampal neurons](#)
- [AXL Mediates Esophageal Adenocarcinoma Cell Invasion through Regulation of Extracellular Acidification and Lysosome Trafficking.](#)