

**Fer Antibody**  
**Purified Mouse Monoclonal Antibody (Mab)**  
**Catalog # AM8466b**

**Specification**

---

**Fer Antibody - Product Information**

Application	IF, WB, FC,E
Primary Accession	<a href="#">P70451</a>
Reactivity	Mouse
Host	Mouse
Clonality	monoclonal
Isotype	IgG2a,k
Calculated MW	94579

**Fer Antibody - Additional Information**

**Gene ID** 14158

**Other Names**

Tyrosine-protein kinase Fer, Proto-oncogene c-Fer, p94-Fer, Fer, Fert2

**Target/Specificity**

This Fer antibody is generated from a mouse immunized with a recombinant protein.

**Dilution**

IF~~1:25

WB~~1:4000

FC~~1:25

**Format**

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Fer Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**Fer Antibody - Protein Information**

**Name** Fer

**Synonyms** Fert2

**Function** Tyrosine-protein kinase that acts downstream of cell surface receptors for growth factors and plays a role in the regulation of the actin cytoskeleton, microtubule assembly,

lamellipodia formation, cell adhesion, cell migration and chemotaxis. Acts downstream of EGFR, KIT, PDGFRA and PDGFRB. Acts downstream of EGFR to promote activation of NF- kappa-B and cell proliferation. May play a role in the regulation of the mitotic cell cycle. Plays a role in the insulin receptor signaling pathway and in activation of phosphatidylinositol 3-kinase. Acts downstream of the activated FCER1 receptor and plays a role in FCER1 (high affinity immunoglobulin epsilon receptor)-mediated signaling in mast cells. Plays a role in the regulation of mast cell degranulation. Plays a role in leukocyte recruitment and diapedesis in response to bacterial lipopolysaccharide (LPS). Phosphorylates CTTN, CTNND1, PTK2/FAK1, GAB1, PECAM1 and PTPN11. May phosphorylate JUP and PTPN1. Can phosphorylate STAT3 according to PubMed:[10878010](#) and PubMed:[19159681](#), but clearly plays a redundant role in STAT3 phosphorylation. According to PubMed:[11134346](#), cells where wild type FER has been replaced by a kinase-dead mutant show no reduction in STAT3 phosphorylation. Phosphorylates TMF1. Isoform 3 lacks kinase activity.

#### Cellular Location

Cytoplasm. Cytoplasm, cytoskeleton. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cell projection. Cell junction. Membrane; Peripheral membrane protein; Cytoplasmic side. Nucleus. Cytoplasm, cell cortex. Note=Detected on microtubules in polarized and motile vascular endothelial cells. Colocalizes with F-actin at the cell cortex. Colocalizes with PECAM1 and CTNND1 at nascent cell-cell contacts (By similarity). Not detected in the nucleus, but detected in the nuclear area surrounding the chromosomes after breakdown of the nuclear envelope during mitosis (PubMed:11339827). {ECO:0000250, ECO:0000269|PubMed:11339827}

#### Tissue Location

Detected in liver and testis. Isoform 4 is detected only in testis (at protein level). Widely expressed


#### Fer Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Fer Antibody - Images






Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized NIH/3T3 (mouse embryonic fibroblast cell line) cells labeling Fer with AM8466b at 1/25 dilution, followed by Dylight® 488-conjugated goat anti-mouse IgG (NA166821) secondary antibody at 1/200 dilution (green). Immunofluorescence image showing cytoplasm staining on NIH/3T3 cell line. The nuclear counter stain is DAPI (blue).



All lanes : Anti-Fer Antibody at 1:4000 dilution Lane 1: NIH/3T3 whole cell lysates Lane 2: mouse testis lysates Lane 3: mouse liver lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 95 kDa Blocking/Dilution buffer: 5% NFDm/TBST.



Overlay histogram showing NIH/3T3 cells stained with AM8466b (green line). The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (AM8466b, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Mouse IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed (NA168821) at 1/400 dilution for 40 min at 37°C. Isotype control antibody (blue line) was mouse IgG1 (1µg/1x10<sup>6</sup> cells) used under the same conditions. Acquisition of >10,000 events was performed.

### Fer Antibody - Background

Tyrosine-protein kinase that acts downstream of cell surface receptors for growth factors and plays a role in the regulation of the actin cytoskeleton, microtubule assembly, lamellipodia formation, cell adhesion, cell migration and chemotaxis. Acts downstream of EGFR, KIT, PDGFRA and PDGFRB. Acts downstream of EGFR to promote activation of NF-kappa-B and cell proliferation. May play a role in the regulation of the mitotic cell cycle. Plays a role in the insulin receptor signaling pathway and in activation of phosphatidylinositol 3-kinase. Acts downstream of the activated FCER1 receptor and plays a role in FCER1 (high affinity immunoglobulin epsilon receptor)-mediated signaling in mast cells. Plays a role in the regulation of mast cell degranulation. Plays a role in leukocyte recruitment and diapedesis in response to bacterial lipopolysaccharide (LPS). Phosphorylates CTTN, CTNND1, PTK2/FAK1, GAB1, PECAM1 and PTPN11. May phosphorylate JUP and PTPN1. Can phosphorylate STAT3 according to PubMed:10878010 and PubMed:19159681, but clearly plays a redundant role in STAT3 phosphorylation. According to PubMed:11134346, cells where wild type FER has been replaced by a kinase-dead mutant show no reduction in STAT3 phosphorylation. Phosphorylates TMF1. Isoform 3 lacks kinase activity.

### Fer Antibody - References

Letwin K., et al. Submitted (OCT-1996) to the EMBL/GenBank/DDBJ databases.  
 Fischman K., et al. Mol. Cell. Biol. 10:146-153(1990).  
 Iwanishi M., et al. J. Biol. Chem. 275:38995-39000(2000).  
 Carninci P., et al. Science 309:1559-1563(2005).  
 Kim L., et al. Mol. Cell. Biol. 15:4553-4561(1995).