

BAP31 Antibody

Purified Mouse Monoclonal Antibody (Mab)
Catalog # AM8468b

Specification

BAP31 Antibody - Product Information

WB, IHC-P,E Application **Primary Accession** P51572 Reactivity Human Host Mouse Clonality monoclonal Isotype IgG1,k 27992 Calculated MW Antigen Region 1-246

BAP31 Antibody - Additional Information

Gene ID 10134

Other Names

B-cell receptor-associated protein 31, BCR-associated protein 31, Bap31, 6C6-AG tumor-associated antigen, Protein CDM, p28, BCAP31, BAP31, DXS1357E

Target/Specificity

This BAP31 antibody is generated from a mouse immunized with a recombinant.

Dilution

WB~~1:2000 IHC-P~~1:25

Format

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

BAP31 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

BAP31 Antibody - Protein Information

Name BCAP31 (<u>HGNC:16695</u>)

Function Functions as a chaperone protein (PubMed:<u>18287538</u>, PubMed:<u>9396746</u>). Is one of the most abundant endoplasmic reticulum (ER) proteins (PubMed:<u>18287538</u>, PubMed:<u>9396746</u>). Plays a role in the export of secreted proteins in the ER, the recognition of abnormally folded protein and



their targeting to the ER associated-degradation (ERAD) (PubMed:18287538, PubMed:9396746). Also serves as a cargo receptor for the export of transmembrane proteins (By similarity). Plays a role in the assembly of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) by stimulating the translocation of NDUFS4 and NDUFB11 from the cytosol to the mitochondria via interaction with TOMM40 (PubMed:31206022). In response to ER stress, delocalizes from the ER-mitochondria contact sites and binds BCL2 (PubMed:31206022). May be involved in CASP8-mediated apoptosis (PubMed:10958671).

Cellular Location

Endoplasmic reticulum membrane; Multi-pass membrane protein Endoplasmic reticulum-Golgi intermediate compartment membrane; Multi-pass membrane protein. Note=May shuttle between the ER and the intermediate compartment/cis-Golgi complex (PubMed:9396746). Associates with the mitochondria-associated endoplasmic reticulum membrane via interaction with TOMM40 (PubMed:31206022)

Tissue Location


Ubiquitous. Highly expressed in neurons and discrete endocrine cells.

BAP31 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

BAP31 Antibody - Images



All lanes : Anti-BAP31 Antibody at 1:2000 dilution Lane 1: A431 whole cell lysates Lane 2: NCCIT whole cell lysates Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 28 kDa Blocking/Dilution buffer: 5% NFDM/TBST.





AM8468b staining BAP31 in human brain sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0. 5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hours at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.

BAP31 Antibody - Background

Functions as a chaperone protein. Is one of the most abundant endoplasmic reticulum (ER) proteins. Plays a role in the export of secreted proteins in the ER, the recognition of abnormally folded protein and their targeting to the ER associated-degradation (ERAD). Also serves as a cargo receptor for the export of transmembrane proteins. May be involved in CASP8- mediated apoptosis.

BAP31 Antibody - References

Mosser J., et al. Genomics 22:469-471(1994). Li E., et al. Eur. J. Biochem. 238:631-638(1996). Adachi T., et al. EMBO J. 15:1534-1541(1996). Ota T., et al. Nat. Genet. 36:40-45(2004). Ross M.T., et al. Nature 434:325-337(2005).