

PPP2R1B Antibody
Purified Mouse Monoclonal Antibody (Mab)
Catalog # AM8469b

Specification

PPP2R1B Antibody - Product Information

Application	IF, WB, IHC-P, FC,E
Primary Accession	P30154
Reactivity	Human, Mouse, Rat
Host	Mouse
Clonality	monoclonal
Isotype	IgG1,K
Calculated MW	66214

PPP2R1B Antibody - Additional Information

Gene ID 5519

Other Names

Serine/threonine-protein phosphatase 2A 65 kDa regulatory subunit A beta isoform, PP2A subunit A isoform PR65-beta, PP2A subunit A isoform R1-beta, PPP2R1B

Target/Specificity

This PPP2R1B antibody is generated from a mouse immunized with a recombinant protein of human PPP2R1B.

Dilution

IF~~1:25
WB~~1:4000
IHC-P~~1:25
FC~~1:25

Format

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PPP2R1B Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

PPP2R1B Antibody - Protein Information

Name PPP2R1B

Function The PR65 subunit of protein phosphatase 2A serves as a scaffolding molecule to


coordinate the assembly of the catalytic subunit and a variable regulatory B subunit.

PPP2R1B Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


PPP2R1B Antibody - Images




Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized U-2 OS (human bone osteosarcoma cell line) cells labeling PPP2R1B with AM8469b at 1/25 dilution, followed by Dylight® 488-conjugated goat anti-mouse IgG (NA166821) secondary antibody at 1/200 dilution (green). Immunofluorescence image showing cytoplasm staining on U-2 OS cell line. Cytoplasmic actin is detected with Dylight® 554 Phalloidin (PD18466410) at 1/100 dilution (red).The nuclear counter stain is DAPI (blue).



All lanes : Anti-PPP2R1B Antibody at 1:4000 dilution Lane 1: Jurkat whole cell lysates Lane 2: human brain lysates Lane 3: mouse brain lysates Lane 4: mouse lung lysates Lane 5: rat brain lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 66 kDa Blocking/Dilution buffer: 5% NFDm/TBST.



AM8469b staining PPP2R1B in human spleen sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hours at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.



Overlay histogram showing Jurkat cells stained with AM8469b (green line). The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (AM8469b, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Mouse IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed(NA168821)) at 1/400 dilution for 40 min at 37°C. Isotype control antibody (blue line) was mouse IgG1 (1µg/1x10⁶ cells) used under the same conditions. Acquisition of >10,000 events was performed.

PPP2R1B Antibody - Background

The PR65 subunit of protein phosphatase 2A serves as a scaffolding molecule to coordinate the assembly of the catalytic subunit and a variable regulatory B subunit.

PPP2R1B Antibody - References

- Baysal B.E., et al. Gene 217:107-116(1998).
- Wang S.S., et al. Science 282:284-287(1998).
- Baysal B.E., et al. Eur. J. Hum. Genet. 9:121-129(2001).
- Ota T., et al. Nat. Genet. 36:40-45(2004).
- Taylor T.D., et al. Nature 440:497-500(2006).