

RAB27B Antibody
Purified Mouse Monoclonal Antibody (Mab)
Catalog # AM8508b

Specification

RAB27B Antibody - Product Information

Application	WB,E
Primary Accession	O00194
Reactivity	Human, Mouse
Host	Mouse
Clonality	monoclonal
Isotype	IgG1,k

RAB27B Antibody - Additional Information

Gene ID 5874

Other Names

Ras-related protein Rab-27B, C25KG, RAB27B

Target/Specificity

This RAB27B antibody is generated from a mouse immunized with a recombinant protein of human RAB27B.

Dilution

WB~~1:2000

Format

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

RAB27B Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

RAB27B Antibody - Protein Information

Name RAB27B

Function Small GTPase which cycles between active GTP-bound and inactive GDP-bound states. In its active state, binds to a variety of effector proteins to regulate homeostasis of late endocytic pathway, including endosomal positioning, maturation and secretion (PubMed:[30771381](#)). Plays a role in NTRK2/TRKB axonal anterograde transport by facilitating the association of NTRK2/TRKB with KLC1 (PubMed:[21775604](#)). May be involved in targeting uroplakins to urothelial apical membranes (By similarity).

Cellular Location

Membrane; Lipid-anchor. Late endosome


Tissue Location

Expressed primarily in testis.

RAB27B Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RAB27B Antibody - Images

Anti-RAB27B Antibody at 1:2000 dilution + mouse stomach lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 25 kDa Blocking/Dilution buffer: 5% NFD/MTBST.

RAB27B Antibody - Background

May be involved in targeting uroplakins to urothelial apical membranes.

RAB27B Antibody - References

- Chen D., et al. *Biochem. Mol. Med.* 60:27-37(1997).
Chen D., et al. Submitted (JUN-1999) to the EMBL/GenBank/DDBJ databases.
Ramalho J.S., et al. *BMC Genet.* 2:2-2(2001).
Puhl H.L. III, et al. Submitted (APR-2002) to the EMBL/GenBank/DDBJ databases.
Ota T., et al. *Nat. Genet.* 36:40-45(2004).