

TERF2IP Antibody
Purified Mouse Monoclonal Antibody (Mab)
Catalog # AM8544b

Specification

TERF2IP Antibody - Product Information

Application	WB, IHC-P,E
Primary Accession	O9NYB0
Reactivity	Human
Host	Mouse
Clonality	monoclonal
Isotype	IgG1,k
Calculated MW	44260

TERF2IP Antibody - Additional Information

Gene ID 54386

Other Names

Telomeric repeat-binding factor 2-interacting protein 1, TERF2-interacting telomeric protein 1, TRF2-interacting telomeric protein 1, Dopamine receptor-interacting protein 5, Repressor/activator protein 1 homolog, RAP1 homolog, hRap1, TERF2IP, DRIP5, RAP1

Target/Specificity

This TE2IP antibody is generated from a mouse immunized with a recombinant protein between 1-399 amino acids from human TE2IP.

Dilution

WB~~1:2000

IHC-P~~1:25

Format

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

TERF2IP Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

TERF2IP Antibody - Protein Information

Name TERF2IP

Synonyms DRIP5, RAP1

Function Acts both as a regulator of telomere function and as a transcription regulator. Involved in the regulation of telomere length and protection as a component of the shelterin complex (telosome). In contrast to other components of the shelterin complex, it is dispensable for telomere capping and does not participate in the protection of telomeres against non-homologous end-joining (NHEJ)- mediated repair. Instead, it is required to negatively regulate telomere recombination and is essential for repressing homology- directed repair (HDR), which can affect telomere length. Does not bind DNA directly: recruited to telomeric double-stranded 5'-TTAGGG-3' repeats via its interaction with TERF2. Independently of its function in telomeres, also acts as a transcription regulator: recruited to extratelomeric 5'-TTAGGG-3' sites via its association with TERF2 or other factors, and regulates gene expression. When cytoplasmic, associates with the I-kappa-B-kinase (IKK) complex and acts as a regulator of the NF-kappa-B signaling by promoting IKK-mediated phosphorylation of RELA/p65, leading to activate expression of NF- kappa-B target genes.

Cellular Location

Nucleus {ECO:0000250|UniProtKB:Q91VL8}. Cytoplasm {ECO:0000250|UniProtKB:Q91VL8}. Chromosome {ECO:0000250|UniProtKB:Q91VL8}. Chromosome, telomere {ECO:0000250|UniProtKB:Q91VL8}. Note=Associates with chromosomes, both at telomeres and in extratelomeric sites. Also exists as a cytoplasmic form, where it associates with the IKK complex {ECO:0000250|UniProtKB:Q91VL8}

Tissue Location

Ubiquitous. Highly expressed.


TERF2IP Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TERF2IP Antibody - Images





All lanes : Anti-TERF2IP Antibody at 1:2000 dilution Lane 1: HepG2 whole cell lysate Lane 2: A431 whole cell lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 44 kDa Blocking/Dilution buffer: 5% NFDN/TBST.



AM8544b staining TE2IP in human testis tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hours at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.

TERF2IP Antibody - Background

Acts both as a regulator of telomere function and as a transcription regulator. Involved in the regulation of telomere length and protection as a component of the shelterin complex (telosome). In contrast to other components of the shelterin complex, it is dispensible for telomere capping and does not participate in the protection of telomeres against non-homologous end-joining (NHEJ)-mediated repair. Instead, it is required to negatively regulate telomere recombination and is essential for repressing homology-directed repair (HDR), which can affect telomere length. Does not bind DNA directly: recruited to telomeric double-stranded 5'-TTAGGG-3' repeats via its interaction with TERF2. Independently of its function in telomeres, also acts as a transcription regulator: recruited to extratelomeric 5'-TTAGGG-3' sites via its association with TERF2 or other factors, and

regulates gene expression. When cytoplasmic, associates with the I-kappa-B-kinase (IKK) complex and acts as a regulator of the NF-kappa-B signaling by promoting IKK-mediated phosphorylation of RELA/p65, leading to activate expression of NF-kappa-B target genes.

TERF2IP Antibody - References

Li B.,et al.Cell 101:471-483(2000).

Lafuente M.J.,et al.Submitted (MAR-2000) to the EMBL/GenBank/DDBJ databases.

Wan D.,et al.Proc. Natl. Acad. Sci. U.S.A. 101:15724-15729(2004).

Ota T.,et al.Nat. Genet. 36:40-45(2004).

Martin J.,et al.Nature 432:988-994(2004).