

MYBBP1A Antibody
Purified Mouse Monoclonal Antibody (Mab)
Catalog # AM8564b

Specification

MYBBP1A Antibody - Product Information

Application	WB,E
Primary Accession	O9BQG0
Reactivity	Human
Host	Mouse
Clonality	monoclonal
Isotype	IgG1,k
Calculated MW	148855

MYBBP1A Antibody - Additional Information

Gene ID 10514

Other Names

Myb-binding protein 1A, MYBBP1A, P160

Target/Specificity

This MYBBP1A antibody is generated from a mouse immunized with a recombinant protein between 1104-1328 amino acids from human MYBBP1A.

Dilution

WB~~1:5000

Format

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

MYBBP1A Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

MYBBP1A Antibody - Protein Information

Name MYBBP1A

Synonyms P160

Function May activate or repress transcription via interactions with sequence specific DNA-binding proteins (By similarity). Repression may be mediated at least in part by histone

deacetylase activity (HDAC activity) (By similarity). Acts as a corepressor and in concert with CRY1, represses the transcription of the core circadian clock component PER2 (By similarity). Preferentially binds to dimethylated histone H3 'Lys-9' (H3K9me2) on the PER2 promoter (By similarity). Has a role in rRNA biogenesis together with PWP1 (PubMed:[29065309](#)).

Cellular Location


Cytoplasm. Nucleus. Nucleus, nucleolus. Note=Shuttles between the nucleus and cytoplasm. Nuclear import may be mediated by KPNA2, while export appears to depend partially on XPO1/CRM1 (By similarity). Predominantly nucleolar. {ECO:0000250|UniProtKB:Q7TPV4}

MYBBP1A Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MYBBP1A Antibody - Images



All lanes : Anti-MYBBP1A Antibody at 1:5000 dilution Lane 1: HeLa whole cell lysate Lane 2: Jurkat whole cell lysate Lane 3: A431 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 149 kDa Blocking/Dilution buffer: 5% NFDN/TBST.

MYBBP1A Antibody - Background

May activate or repress transcription via interactions with sequence specific DNA-binding proteins. Repression may be mediated at least in part by histone deacetylase activity (HDAC activity). Acts as a corepressor and in concert with CRY1, represses the transcription of the core circadian clock component PER2. Preferentially binds to dimethylated histone H3 'Lys-9' (H3K9me2) on the PER2 promoter.

MYBBP1A Antibody - References

Keough R., et al. Genomics 62:483-489(1999).
Wiemann S., et al. Genome Res. 11:422-435(2001).
Bechtel S., et al. BMC Genomics 8:399-399(2007).
Scherl A., et al. Mol. Biol. Cell 13:4100-4109(2002).
Olsen J.V., et al. Cell 127:635-648(2006).