

**MYC Antibody**  
**Purified Mouse Monoclonal Antibody (Mab)**  
**Catalog # AM8600b**

**Specification**

---

**MYC Antibody - Product Information**

Application	<b>WB,E</b>
Primary Accession	<a href="#">P01106</a>
Reactivity	<b>Human</b>
Host	<b>Mouse</b>
Clonality	<b>monoclonal</b>
Isotype	<b>IgG1,k</b>
Calculated MW	<b>50565</b>

**MYC Antibody - Additional Information**

**Gene ID** 4609

**Other Names**

Myc proto-oncogene protein, Class E basic helix-loop-helix protein 39, bHLHe39, Proto-oncogene c-Myc, Transcription factor p64, MYC, BHLHE39

**Target/Specificity**

This MYC antibody is generated from a mouse immunized with a recombinant protein of human MYC Isform 2.

**Dilution**

WB~~1:2000

**Format**

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

MYC Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**MYC Antibody - Protein Information**

**Name** MYC

**Synonyms** BHLHE39

**Function** Transcription factor that binds DNA in a non-specific manner, yet also specifically recognizes the core sequence 5'-CAC[GA]TG-3' (PubMed:[24940000](#), PubMed:[25956029](#)). Activates

the transcription of growth-related genes (PubMed:[24940000](#), PubMed:[25956029](#)). Binds to the VEGFA promoter, promoting VEGFA production and subsequent sprouting angiogenesis (PubMed:[24940000](#), PubMed:[25956029](#)). Regulator of somatic reprogramming, controls self-renewal of embryonic stem cells (By similarity). Functions with TAF6L to activate target gene expression through RNA polymerase II pause release (By similarity). Positively regulates transcription of HNRNPA1, HNRNPA2 and PTBP1 which in turn regulate splicing of pyruvate kinase PKM by binding repressively to sequences flanking PKM exon 9, inhibiting exon 9 inclusion and resulting in exon 10 inclusion and production of the PKM M2 isoform (PubMed:[20010808](#)).

#### Cellular Location


Nucleus, nucleoplasm. Nucleus, nucleolus. Nucleus. Cytoplasm Note=Localization to the nucleolus is dependent on HEATR1

#### MYC Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### MYC Antibody - Images



All lanes : Anti-MYC Antibody at 1:2000 dilution Lane 1: HeLa whole cell lysate Lane 2: K562 whole cell lysate Lane 3: Jurkat whole cell lysate Lane 4: Raji whole cell lysate Lane 5: LNCaP whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 51 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

#### MYC Antibody - Background

Transcription factor that binds DNA in a non-specific manner, yet also specifically recognizes the

core sequence 5'- CAC[GA]TG-3'. Activates the transcription of growth-related genes.

### **MYC Antibody - References**

- Batley J., et al. Cell 34:779-787(1983).  
Bernard O., et al. EMBO J. 2:2375-2383(1983).  
Colby W.W., et al. Nature 301:722-725(1983).  
Watt R., et al. Nature 303:725-728(1983).  
Rabbitts T.H., et al. Nature 306:760-765(1983).