

**ADK Antibody (N-term)**  
**Purified Mouse Monoclonal Antibody (Mab)**  
**Catalog # AM8619b**

**Specification**

---

**ADK Antibody (N-term) - Product Information**

Application	<b>WB, IHC-P,E</b>
Primary Accession	<a href="#">P55263</a>
Reactivity	<b>Human</b>
Host	<b>Mouse</b>
Clonality	<b>monoclonal</b>
Isotype	<b>IgG1,k</b>
Calculated MW	<b>40545</b>

**ADK Antibody (N-term) - Additional Information**

**Gene ID** 132

**Other Names**

Adenosine kinase, AK, 2.7.1.20, Adenosine 5'-phosphotransferase, ADK

**Target/Specificity**

This ADK antibody is generated from a mouse immunized with a recombinant protein between 1-345 amino acids from human ADK.

**Dilution**

WB~~1:4000

IHC-P~~1:25

**Format**

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

ADK Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**ADK Antibody (N-term) - Protein Information**

**Name** ADK ([HGNC:257](#))

**Function** Catalyzes the phosphorylation of the purine nucleoside adenosine at the 5' position in an ATP-dependent manner. Serves as a potential regulator of concentrations of extracellular adenosine and intracellular adenine nucleotides.

**Cellular Location**

[Isoform 1]: Nucleus


**Tissue Location**

Widely expressed. Highest level in placenta, liver, muscle and kidney.


**ADK Antibody (N-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**ADK Antibody (N-term) - Images**

All lanes : Anti-ADK Antibody (N-term) at 1:4000 dilution Lane 1: HepG2 whole cell lysate Lane 2: HL-60 whole cell lysate Lane 3: Human liver lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 45 kDa Blocking/Dilution buffer: 5% NFDm/TBST.



AM8619b staining ADK in human kidney tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hour at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.

#### **ADK Antibody (N-term) - Background**

ATP dependent phosphorylation of adenosine and other related nucleoside analogs to monophosphate derivatives. Serves as a potential regulator of concentrations of extracellular adenosine and intracellular adenine nucleotides.

#### **ADK Antibody (N-term) - References**

- Spychala J., et al. Proc. Natl. Acad. Sci. U.S.A. 93:1232-1237(1996).  
Singh B., et al. Eur. J. Biochem. 241:564-571(1996).  
McNally T., et al. Biochem. Biophys. Res. Commun. 231:645-650(1997).  
Ota T., et al. Nat. Genet. 36:40-45(2004).  
Deloukas P., et al. Nature 429:375-381(2004).