

SNCA Antibody

Purified Mouse Monoclonal Antibody (Mab)
Catalog # AM8649b

Specification

SNCA Antibody - Product Information

Application
Primary Accession
Other Accession
Reactivity
Host

Clonality Isotype Calculated

Calculated MW

SNCA Antibody - Additional Information

Gene ID 6622

Other Names

Alpha-synuclein, Non-A beta component of AD amyloid, Non-A4 component of amyloid precursor, NACP, SNCA, NACP, PARK1

WB,E

P37840

Mouse

IgG1, ĸ

14460

P61140, P61144, P61145

Human, Mouse

monoclonal

Target/Specificity

This antibody is generated from a mouse immunized a recombinant protein from human.

Dilution

WB~~1:2000

Format

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

SNCA Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

SNCA Antibody - Protein Information

Name SNCA

Synonyms NACP, PARK1

Function Neuronal protein that plays several roles in synaptic activity such as regulation of synaptic vesicle trafficking and subsequent neurotransmitter release (PubMed: 20798282,



PubMed: 26442590, PubMed: 28288128, PubMed: 30404828). Participates as a monomer in synaptic vesicle exocytosis by enhancing vesicle priming, fusion and dilation of exocytotic fusion pores (PubMed: 28288128, PubMed: 30404828). Mechanistically, acts by increasing local Ca(2+) release from microdomains which is essential for the enhancement of ATP-induced exocytosis (PubMed: 30404828). Acts also as a molecular chaperone in its multimeric membrane-bound state, assisting in the folding of synaptic fusion components called SNAREs (Soluble NSF Attachment Protein REceptors) at presynaptic plasma membrane in conjunction with cysteine string protein-alpha/DNAJC5 (PubMed: 20798282). This chaperone activity is important to sustain normal SNARE-complex assembly during aging (PubMed: 20798282). Also plays a role in the regulation of the dopamine neurotransmission by associating with the dopamine transporter (DAT1) and thereby modulating its activity (PubMed: 26442590).

Cellular Location

Cytoplasm. Membrane Nucleus Synapse. Secreted. Cell projection, axon {ECO:0000250|UniProtKB:O55042}. Note=Membrane-bound in dopaminergic neurons (PubMed:15282274). Expressed and colocalized with SEPTIN4 in dopaminergic axon terminals, especially at the varicosities (By similarity). {ECO:0000250|UniProtKB:O55042, ECO:0000269|PubMed:15282274}

Tissue Location


Highly expressed in presynaptic terminals in the central nervous system. Expressed principally in brain

SNCA Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

SNCA Antibody - Images





All lanes: Anti-SNCA at 1:2000 dilution Lane 1: Human brain lysate Lane 2: Mouse brain lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 14 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

SNCA Antibody - Background

May be involved in the regulation of dopamine release and transport. Induces fibrillization of microtubule-associated protein tau. Reduces neuronal responsiveness to various apoptotic stimuli, leading to a decreased caspase-3 activation.

SNCA Antibody - References

Ueda K.,et al.Proc. Natl. Acad. Sci. U.S.A. 90:11282-11286(1993). Campion D.,et al.Genomics 26:254-257(1995). Ueda K.,et al.Biochem. Biophys. Res. Commun. 205:1366-1372(1994). Xia Y.,et al.Submitted (JAN-1996) to the EMBL/GenBank/DDBJ databases. Touchman J.W.,et al.Genome Res. 11:78-86(2001).