

**CD276 Antibody**  
**Purified Mouse Monoclonal Antibody (Mab)**  
**Catalog # AM8677b**

**Specification**

---

**CD276 Antibody - Product Information**

Application	WB, IHC, IHC-P,E
Primary Accession	<a href="#">Q5ZPR3</a>
Reactivity	Human
Predicted	Human
Host	Mouse
Clonality	monoclonal
Isotype	IgG1, $\kappa$

**CD276 Antibody - Additional Information**

**Gene ID** 80381

**Other Names**

CD276 antigen, 4Ig-B7-H3, B7 homolog 3, B7-H3, Costimulatory molecule, CD276, CD276, B7H3

**Target/Specificity**

This CD276 antibody is generated from a mouse immunized with a recombinant protein from the human region of human CD276.

**Dilution**

WB~~1:4000

IHC~~1:400

IHC-P~~1:25

**Format**

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

CD276 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**CD276 Antibody - Protein Information**

**Name** CD276

**Synonyms** B7H3

**Function** May participate in the regulation of T-cell-mediated immune response. May play a

protective role in tumor cells by inhibiting natural-killer mediated cell lysis as well as a role of marker for detection of neuroblastoma cells. May be involved in the development of acute and chronic transplant rejection and in the regulation of lymphocytic activity at mucosal surfaces. Could also play a key role in providing the placenta and fetus with a suitable immunological environment throughout pregnancy. Both isoform 1 and isoform 2 appear to be redundant in their ability to modulate CD4 T-cell responses. Isoform 2 is shown to enhance the induction of cytotoxic T-cells and selectively stimulates interferon gamma production in the presence of T-cell receptor signaling.

#### Cellular Location

Membrane; Single-pass type I membrane protein

#### Tissue Location


Ubiquitous but not detectable in peripheral blood lymphocytes or granulocytes. Weakly expressed in resting monocytes Expressed in dendritic cells derived from monocytes. Expressed in epithelial cells of sinonasal tissue. Expressed in extravillous trophoblast cells and Hofbauer cells of the first trimester placenta and term placenta.

### CD276 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### CD276 Antibody - Images




All lanes : Anti-CD276 Antibody at 1:4000 dilution Lane 1: 293T-17 whole cell lysate Lane 2: A-431 whole cell lysate Lane 3: U-20S whole cell lysate Lane 4: LNCap whole cell lysate Lane 5: HeLa whole cell lysate Lane 6: MCF-7 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted


band size : 57 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Immunohistochemical analysis of paraffin-embedded Human tonsil section using CD276 Antibody (Cat#AM8677B). AM8677B was diluted at 1:400 dilution. A undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB staining.



Immunohistochemical analysis of CD276 Antibody in 293T(untransfected) cell sections. Cell was fixed with formaldehyde and blocked with super block for 10 minutes at room temperature; antigen retrieval was by heat mediation with a EDTA buffer (pH9. 0). Samples were incubated with primary antibody (1:500) for 1 hours at room temperature. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.



Immunohistochemical analysis of CD276 Antibody in MCF-7 cell sections. Cell was fixed with formaldehyde and blocked with super block for 10 minutes at room temperature; antigen retrieval was by heat mediation with a EDTA buffer (pH9. 0). Samples were incubated with primary antibody (1:25) for 1 hours at room temperature. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.

### **CD276 Antibody - Background**

May participate in the regulation of T-cell-mediated immune response. May play a protective role in tumor cells by inhibiting natural-killer mediated cell lysis as well as a role of marker for detection of neuroblastoma cells. May be involved in the development of acute and chronic transplant rejection and in the regulation of lymphocytic activity at mucosal surfaces. Could also play a key role in providing the placenta and fetus with a suitable immunological environment throughout pregnancy. Both isoform 1 and isoform 2 appear to be redundant in their ability to modulate CD4 T-cell responses. Isoform 2 is shown to enhance the induction of cytotoxic T-cells and selectively stimulates interferon gamma production in the presence of T-cell receptor signaling.

### **CD276 Antibody - References**

Chapoval A.I.,et al.Nat. Immunol. 2:269-274(2001).  
Steinberger P.,et al.J. Immunol. 172:2352-2359(2004).  
Clark H.F.,et al.Genome Res. 13:2265-2270(2003).  
Ota T.,et al.Nat. Genet. 36:40-45(2004).  
Otsuki T.,et al.DNA Res. 12:117-126(2005).