

RAB5B Antibody
Purified Mouse Monoclonal Antibody (Mab)
Catalog # AM8689b

Specification

RAB5B Antibody - Product Information

Application	WB,E
Primary Accession	P61020
Other Accession	P61021 , Q5RBG1
Reactivity	Human, Mouse
Predicted	Mouse
Host	Mouse
Clonality	monoclonal
Isotype	IgG1,k
Calculated MW	23707

RAB5B Antibody - Additional Information

Gene ID 5869

Other Names

Ras-related protein Rab-5B, RAB5B

Target/Specificity

This RAB5B antibody is generated from a mouse immunized with a recombinant protein of human RAB5B.

Dilution

WB~~1:2000

Format

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

RAB5B Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

RAB5B Antibody - Protein Information

Name RAB5B

Function Protein transport. Probably involved in vesicular traffic.

Cellular Location


Cell membrane; Lipid-anchor; Cytoplasmic side. Early endosome membrane; Lipid-anchor. Melanosome. Note=Enriched in stage I melanosomes

RAB5B Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RAB5B Antibody - Images



Anti-RAB5B Antibody at 1:2000 dilution + A431 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 24 kDa Blocking/Dilution buffer: 5% NFDN/TBST.



All lanes : Anti-RAB5B Antibody at 1:2000 dilution Lane 1: HeLa whole cell lysate Lane 2: HepG2 whole cell lysate Lane 3: Jurkat whole cell lysate Lane 4: NIH-3T3 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 24 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

RAB5B Antibody - Background

Protein transport. Probably involved in vesicular traffic (By similarity).

RAB5B Antibody - References

- Wilson D.B.,et al.J. Clin. Invest. 89:996-1005(1992).
- Puhl H.L. III,et al.Submitted (APR-2002) to the EMBL/GenBank/DDBJ databases.
- Ota T.,et al.Nat. Genet. 36:40-45(2004).
- Bechtel S.,et al.BMC Genomics 8:399-399(2007).
- Scherer S.E.,et al.Nature 440:346-351(2006).