

PNOC Antibody (Center)
Purified Mouse Monoclonal Antibody (Mab)
Catalog # AM8712b

Specification

PNOC Antibody (Center) - Product Information

Application	WB,E
Primary Accession	O13519
Reactivity	Human
Predicted	Human
Host	Mouse
Clonality	monoclonal
Isotype	IgG2b, κ
Calculated MW	20295

PNOC Antibody (Center) - Additional Information

Gene ID 5368

Other Names

Prepronociceptin, Nocistatin, Nociceptin, Orphanin FQ, PPNOC, Orphanin FQ2, PNOC, OFQ

Target/Specificity

This PNOC antibody is generated from a mouse immunized with a recombinant protein from human PNOC.

Dilution

WB~~1:8000

Format

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PNOC Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

PNOC Antibody (Center) - Protein Information

Name PNOC

Synonyms OFQ

Function [Nociceptin]: Ligand of the opioid receptor-like receptor OPRL1. It may act as a

transmitter in the brain by modulating nociceptive and locomotor behavior. May be involved in neuronal differentiation and development.

Cellular Location

Secreted.

Tissue Location


Predominantly expressed in the brain and spinal cord. Also expressed and secreted by peripheral blood neutrophils following degranulation.

PNOC Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PNOC Antibody (Center) - Images



Anti-PNOC Antibody (Center) at 1:8000 dilution + PNOC recombinant protein Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 18 kDa Blocking/Dilution buffer: 5% NFD/MTBST.

PNOC Antibody (Center) - Background

Nociceptin: Ligand of the opioid receptor-like receptor OPRL1. It may act as a transmitter in the brain by modulating nociceptive and locomotor behavior. May be involved in neuronal differentiation and development.

PNOC Antibody (Center) - References

Mollereau C.,et al.Proc. Natl. Acad. Sci. U.S.A. 93:8666-8670(1996).
Nothacker H.-P.,et al.Proc. Natl. Acad. Sci. U.S.A. 93:8677-8682(1996).
Fiset M.E.,et al.Biochemistry 42:10498-10505(2003).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Ebert L.,et al.Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases.