

**Phospho-Tyr1336 NMDA Receptor NR2B Subunit Antibody**  
Affinity purified rabbit polyclonal antibody  
Catalog # AN1094

## Specification

---

### Phospho-Tyr1336 NMDA Receptor NR2B Subunit Antibody - Product Information

Application	WB, IHC
Primary Accession	<a href="#">Q00960</a>
Reactivity	Rat
Predicted	Human, Mouse, Monkey
Host	Rabbit
Clonality	polyclonal
Calculated MW	180 KDa

### Phospho-Tyr1336 NMDA Receptor NR2B Subunit Antibody - Additional Information

Gene ID	24410
Gene Name	GRIN2B

#### Other Names

Glutamate receptor ionotropic, NMDA 2B, GluN2B, Glutamate [NMDA] receptor subunit epsilon-2, N-methyl D-aspartate receptor subtype 2B, NMDAR2B, NR2B, Grin2b

#### Target/Specificity

Synthetic phospho-peptide corresponding to amino acid residues surrounding Tyr1336 conjugated to KLH.

#### Dilution

WB~~ 1:1000

IHC~~ 1:400

#### Format

Prepared from rabbit serum by affinity purification via sequential chromatography on phospho- and dephosphopeptide affinity columns.

#### Antibody Specificity

Specific for ~180k NMDAR NR2B subunit protein phosphorylated at Tyr1336. Immunolabeling of the NMDAR NR2B subunit band is blocked by  $\lambda$ -phosphatase treatment.

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### Precautions

Phospho-Tyr1336 NMDA Receptor NR2B Subunit Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

#### Shipping


Blue Ice

## Phospho-Tyr1336 NMDA Receptor NR2B Subunit Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Phospho-Tyr1336 NMDA Receptor NR2B Subunit Antibody - Images



Western blot of rat hippocampal lysate showing specific immunolabeling of the ~180k NR2B subunit phosphorylated at Tyr1336 (Control). Phosphospecificity is shown in the second lane (lambda-phosphatase: λ-Ptase). The blot is identical to the control except that it was incubated in λ-Ptase (1200 units for 30 min) before being exposed to the phospho-Tyr1336 NMDA NR2B antibody. The immunolabeling is completely eliminated by treatment with λ-Ptase.



Immunostaining of mouse dentate gyrus 48 hr post TMT treatment showing NR2B when phosphorylated at Tyr1336 in green and nuclei in blue.

## Phospho-Tyr1336 NMDA Receptor NR2B Subunit Antibody - Background

The NMDAR plays an essential role in memory, neuronal development and it has also been implicated in several disorders of the central nervous system including Alzheimer's, epilepsy and ischemic neuronal cell death (Grosshans et al., 2002; Wenthold et al., 2003; Carroll and Zukin, 2002). The rat NMDAR1 (NR1) was the first subunit of the NMDAR to be cloned. The NR1 protein can form NMDA activated channels when expressed in *Xenopus* oocytes but the currents in such channels are much smaller than those seen in situ. Channels with more physiological characteristics are produced when the NR1 subunit is combined with one or more of the NMDAR2 (NR2 A-D) subunits (Ishii et al., 1993). Phosphorylation of Tyr1336 is thought to potentiate NMDA receptor-dependent influx of calcium (Takasu et al., 2002) and ischemia may also increase the phosphorylation of this site (Takagi et al., 2003).

### **Phospho-Tyr1336 NMDA Receptor NR2B Subunit Antibody - References**

Carroll RC, Zukin RS (2002) NMDA-receptor trafficking and targeting: implications for synaptic transmission and plasticity. *Trends Neurosci* 25:571-577.

Grosshans DR, Clayton DA, Coultrap SJ, Browning MD (2002) LTP leads to rapid surface expression of NMDA but not AMPA receptors in adult rat CA1. *Nat Neurosci* 5:27-33.

Ishii T, Moriyoshi K, Sugihara H, Sakurada K, Kadotani H, Yokoi M, Akazawa C, Shigemoto R, Mizuno N, Masu M, Nakanishi S (1993) Molecular characterization of the family of the N-methyl- D-aspartate receptor subunits. *J Biol Chem* 268:2836-2843.

Takasu, MA, Dalva, MB, Zigmond, RE, Greenberg, ME (2002) Modulation of NMDA Receptor -Dependent Calcium Influx and Gene Expression Through EphB Receptors. *Science* 295:491-495.

Wenthold RJ, Prybylowski K, Standley S, Sans N, Petralia RS (2003) Trafficking of NMDA receptors. *Annu Rev Pharmacol Toxicol* 43:335-358.

Tatyana Chernova, Joern R. Steinert, Christopher J. Guerin, Pierluigi Nicotera, Ian D. Forsythe, and Andrew G. Smith (2007) Neurite Degeneration Induced by Heme Deficiency Mediated via Inhibition of NMDA Receptor-Dependent Extracellular Signal-Regulated Kinase 1/2 Activation *J. Neurosci.*, 27: 8475 - 8485.