

Synaptotagmin Antibody
Affinity purified rabbit polyclonal antibody
Catalog # AN1109

Specification

Synaptotagmin Antibody - Product Information

| | |
|-------------------|---------------------------------------|
| Application | WB |
| Primary Accession | P21707 |
| Reactivity | Human, Mouse, Rat |
| Predicted | Bovine, Chicken, Human, Mouse, Monkey |
| Host | Rabbit |
| Clonality | polyclonal |
| Calculated MW | 65 KDa |

Synaptotagmin Antibody - Additional Information

| | |
|--------------------|---|
| Gene ID | 25716 |
| Gene Name | SYT1 |
| Other Names | Synaptotagmin-1, Synaptotagmin I, SytI, p65, Syt1 |

Target/Specificity

Synthetic peptide corresponding to amino acid residues specific to synaptotagmin 1 conjugated to KLH.

Dilution

WB~~ 1:1000

Format

Prepared from rabbit serum by affinity purification using a column to which the peptide immunogen was coupled

Antibody Specificity

Specific for ~65k synaptotagmin I protein

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Synaptotagmin Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Shipping


Blue Ice

Synaptotagmin Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Synaptotagmin Antibody - Images




All lanes : Anti-Syt1 Antibody at 1:1000 dilution Lane 1: mouse brain lysate Lane 2: rat brain lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 48 kDa Blocking/Dilution buffer: 5% NFDm/TBST.



Western blot analysis of lysates from rat brain, mouse brain, human brain tissue lysate(from left to right), using Syt1 Antibody(Cat. #AN1109). AN1109 was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysates at 35ug

per lane.



Western blot of rat cortex lysate showing specific labeling of the ~65k synaptotagmin protein.

Synaptotagmin Antibody - Background

Synaptotagmin 1 is a synaptic vesicle membrane glycoprotein that is widely expressed throughout the CNS and is generally thought to act as the Ca^{2+} sensor in the regulation of exocytosis and neurotransmitter release (Littleton and Bellen 1995). Recent studies indicate that synaptotagmin's Ca^{2+} mediated binding of SNAP25 is essential for the Ca^{2+} dependent triggering of membrane fusion (Zhang et al., 2002). It has recently been demonstrated that discrete residues within the c(2)b binding domain of synaptotagmin 1 independently specify endocytic rate and synaptic vesicle size (Poskanzer et al., 2006).

Synaptotagmin Antibody - References

Littleton JT, Bellen HJ (1995) Synaptotagmin controls and modulates synaptic vesicle fusion in a Ca^{2+} dependent manner. *Trends Neurosci.* Apr;18(4):177-83.
Zhang X, Kim-Miller MJ, Fukuda M, Kowalchuk JA, Martin TF (2002) Ca^{2+} dependent synaptotagmin binding to SNAP25 is essential for Ca^{2+} triggered exocytosis. *Neuron* May 16;34(4):599-611.
Poskanzer KE, Fetter RD, Davis GW (2006) Discrete residues in the c(2)b domain of synaptotagmin 1 independently specify endocytic rate and synaptic vesicle size. *Neuron* Apr 6;50(1):49-62.